


2.49

LAWN FUNGICIDE

For Control and Prevention of Dollar and Copper Spot, Helminthosporium-Curvularia (Fading or Melting Out) and Brown Patch in Lawn Grasses.

COVERS 8000 SQ. FT.
SINCE 1855



Use with Well-Lawn Fungicide when the weather prevails and diseases which occur from May to June to Mid-August may be controlled by spraying with 2 1/2 tablespoons of Fungicide in 3 to 5 gals. of water per 1000 sq. ft. NOTE: The amount of water used is not critical as long as proper amount of fungicide is applied. If pressure type sprayer (type carried on back) is used wear an appropriate respirator mask. Repeated applications are necessary for disease control which is usually obtained after 3 applications. There is no noticeable control after one treatment. Applications should be from 7 to 10 days apart.

DIRECTIONS: Diseases which occur from May to June to Mid-August may be controlled by spraying with 2 1/2 tablespoons of Fungicide in 3 to 5 gals. of water per 1000 sq. ft. NOTE: The amount of water used is not critical as long as proper amount of fungicide is applied. If pressure type sprayer (type carried on back) is used wear an appropriate respirator mask. Repeated applications are necessary for disease control which is usually obtained after 3 applications. There is no noticeable control after one treatment. Applications should be from 7 to 10 days apart.

IMPORTANT: 1. Normal rainfall following application will not effect results. 2. Keep lawn mowed at 1 to 1 1/2 inches during treatment. 3. Well-Lawn Fungicide will prevent and control diseases only if it works to kill or effect weeds. 4. Spraying equipment should be cleaned after use.

WARNING: Wash thoroughly after handling and before smoking or eating. May be fatal if swallowed. Do not apply or allow to drift in areas occupied by unprotected persons or beneficial animals. Do not breathe spray mist. If pressure type sprayer (type carried on back) is used wear an appropriate respirator.

EXTERNAL: If spilled on skin wash off immediately with warm water and soap. **INTERNAL:** If accidentally swallowed give raw white of egg. Induce vomiting with mixture of salt, soap or mustard. Call a doctor immediately.

NOTE: Follow directions carefully as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all liability. Directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS: Cadmium Chloride 12.3%
INERT INGREDIENTS: 87.7%

NET CONTENTS 8 OUNCES

12/67



2.49

Gro-Well Lawn Fungicide will control and prevent turf diseases in Kentucky Blue, Merion, Fescue, Bent & Zoysia lawns. It may be applied without fear of injury or discoloration to newly seeded areas as well as established lawns with a Gro-Well Lawn Sprayer.

DIRECTIONS: Diseases which occur during April and May when wet cool weather prevails and diseases which occur during hot summer months from Mid-June to Mid-August may be controlled by using 1 oz. (2 Tablespoons) of Fungicide in 3 to 5 gals. of water per 1000 sq. ft. **NOTE:** The amount of water used is not critical as long as proper amount of fungicide is applied. If pressure type sprayer (type carried on back) is used wear an appropriate respirator mask. Repeated applications are necessary for disease control which is usually obtained after 3 applications. There is no noticeable control after one treatment. Applications should be from 7 to 10 days apart.

IMPORTANT: 1. Normal rainfall following application will not effect results. 2. Keep lawn mowed at 1 to 1½ inches during treatment. 3. Gro-Well Lawn Fungicide will prevent and control diseases only . . . it will not kill or effect weeds. 4. Spraying equipment should be cleaned after use.

WARNING: Wash thoroughly after handling and before smoking or eating. May be fatal if swallowed. Do not apply or allow to drift in areas occupied by unprotected persons or beneficial animals. Do not breathe spray mist. If pressure type sprayer (type carried on back) is used, wear an appropriate respirator.

EXTERNAL: If spilled on skin wash off immediately with warm water and soap. **INTERNAL:** If accidentally swallowed, give raw white of egg. Induce vomiting with mixture of salt, soap or mustard. Call a doctor immediately.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS: Cadmium Chloride 20.1%
INERT INGREDIENTS: 79.9%

Total Cadmium (Cd) 12.3% **TOTAL** 100.0%
Sold by: **NET CONTENTS 8 OUNCES** **USDA Reg. No 5535-2**
 J. & L. ADIKES, INC., JAMAICA, NEW YORK

INERT INGREDIENTS FOR USDA Reg. No. 5535-2

Ammonium Nitrate	2.0%
Water	<u>77.9%</u>
Total	79.9%

ACCEPTED

JUN 25 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 2332-2 SUBJECT
TO ATTACHED COMMENTS.