

FRONT PANEL

Since 1855



FOR FLOWERS, FRUIT TREES, SHRUBS,
VEGETABLES AND GENERAL OUTDOOR SPRAYING

FOR INSECTS & SCALES

MAKES UP TO 100 GALLONS OF SPRAY

CAUTION:

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

[283]

50% MALATHION SPRAY

GRO-WELL MALATHION may be used to control practically every major pest attacking flowers, shrubs, trees, garden vegetables and fruits. Read cautions carefully before using. A sprayer which attaches to hose, such as the GRO-WELL Shrub and Tree Sprayer may be used.

GENERAL DIRECTIONS

Use the dilution rate given below for insects and plants listed mixing with 1 gallon of water. When spraying, wet all plant surfaces, paying particular attention to the undersides of leaves, as this is where most insects feed. On all uses repeat as necessary or at 2 week intervals. (See note below on vegetables and fruit. On fruits apply at petal fall and every 10 to 14 days until control is achieved.) Do not store diluted material. Make new dilution for each use.

INSECTS	FOR USE ON	DOSAGE
<u>GROUP 1</u> Aphids, Birch Leaf Miner, Bugs (Lace & Mealy), Japanese Beetle, Thrips, White fly.	Andromedas, Arborvitae, Asters, Azaleas, Birch, Boxwood, Camelias, Carnation, Dahlias, Elm, Euonymus, Fruit Trees (Apple, Grapes, Peaches, Pears), Gardenias, Gladiolus, Hibiscus, Holly, Hydrangeas, Junipers, Lilac, Magnolia, Maple, Mums, Oak, Pine, Rhododendrons, Roses, Sycamore, Taxus, vegetables (Beans, Cabbage, Corn, Cucumbers, Lettuce, Peas, Peppers, Tomatoes), All ornamentals and Shade Trees.	<u>GROUP 1</u> One teaspoonful in 1 gallon of water.
<u>GROUP 2</u> Boxwood Leaf Miner, Oak Kermes, Red Spider Mites, Scale (Crawlers), Tent Caterpillar.	ACCEPTED JUN 30-1968	<u>GROUP 2</u> Two teaspoonfuls in a gallon of water.
<u>GROUP 3</u> Flies, Mosquitoes	Outside walls, fences, patios, garbage containers.	<u>GROUP 3</u> 5 tablespoonfuls (2½ oz.) in one gallon of water
<u>GROUP 4</u> Codling Moth, Oriental Fruit Moth, Plum Curculio	On Fruit Trees listed above.	<u>GROUP 4</u> Two teaspoonfuls in a gallon of water every 10 to 14 days.

NOTE: Do not spray fruits or vegetables named above later than 10 days before harvest.
 Do not use on Boston, Maidenhair, Pteris and Crassula Ferns.
 Do not use on Japanese Red Maple.

[383]

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid breathing spray mist and contact with skin, eyes and clothing. If gotten into eyes, flush profusely with water and contact doctor. For skin contact wash thoroughly with water and soap, also before eating or smoking. Do not apply to humans or animals. Do not apply or allow to drift into areas occupied by unprotected persons or animals. Keep away from open flame, non-inflammable when diluted as directed. Avoid contamination of feed and foodstuffs. Do not store diluted material, make a new dilution for each spraying job. If spilled, flush area with plenty of clear water. Dispose of this bottle when empty.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

NOTE: Follow directions carefully. As use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS:

*Malathion	50.00%
Xylene	42.40%

INERT INGREDIENTS:	7.60%
TOTAL	100.00%

*O,O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate.

NOTE TO PHYSICIAN:

Malathion upon repeated prolonged careless use, may cause cholinesterase inhibition. Atropine is antidotal.

SOLD BY

J. & L. ADIKES, INC. JAMAICA, NEW YORK

NET CONTENTS - 1 QUART (32 ozs.)

U.S.D.A. Reg. No. 5535-6

July 30-1963
55356