


FRONT
PANEL

1.49

FRUIT TREE *Spray*

MAKES 10 GALLONS
OF SPRAY

SINCE 1855



INSECTICIDE
FUNGICIDE
MITICIDE

CONTAINS
MALATHION
CAPTAN
SEVIN

ACCEPTED

NEW 216

U. S. DEPARTMENT OF AGRICULTURE
FOR ECONOMIC CONTROL ACT
ED UNDER NO. 5535-7 SUBJECT
TO ATTACHED COMMENTS.

REAR
PANEL

GRO-WELL FRUIT TREE SPRAY is a combination insecticide and fungicide for the control of insects and diseases on APPLE, PEACH, CHERRY, PEAR and certain other fruit trees.

GRO-WELL FRUIT TREE SPRAY may be used to control: Green Apple Aphids, Woolly Apple Aphid, Codling Moth, Apple Maggot, European Red Mites, Cleyer Mites, Two Spotted Mites, Willamette Mites, Eye Spotted Bud Moti, Fruit Tree Leaf holler, Apple Red Bug, Pear Psylla, Pear Midge, Pear Plant Bug, Tarnished Plant Bug on APPLES and PEARS; Oriental Fruit Moth, Plum Curculio, Peach Tree Borer, Cottony Peach Scale, Japanese Beetle on PEACHES; European Red Mite on PLUMS and PRUNES; Black Cherry Aphid, Two Spotted Mites and Tent Caterpillar on CHERRIES; Apple Scab, Brown Rot, Leaf Spot on APPLES, PEACHES, PLUMS, PRUNES and CHERRIES; Cherry Fruit Ely, Cherry Fruitworm and Mealy Plum Aphid on CHERRIES, PLUMS and PRUNES; Grape Leafoppers, Grape Leaf Skeletonizer, Grape Berry Moth, Japanese Beetles, Red Banded Leaf Roller and Downey Mildew on Grapes; Spittlebug, Leaf Roller and Strawberry Weevil on Strawberries. Blueberry Maggot, Cherry Fruitworm and Japanese Beetles on Blueberries

NOTE: For Peach Tree Borer, make two sprays starting about July 7-15 and another about August 1-10. Consult Extension Authorities for exact timing in your area. Where disease control close to harvest is necessary, apply a suitable fungicide alone in a manner which will not leave residues in excess of established tolerances. Where mite control is a serious problem add 1 tablespoonful of GRO-WELL KELTHANE per gallon of spray mixture.

DIRECTIONS: SPRAYING—Add the required amount of material to water and stir thoroughly. Pour the mixture into the sprayer and agitate while spraying in order to obtain a thorough and even coverage. In a small sprayer, shake often for agitation. A sprayer that attaches to the hose such as the Gro-Well Shrub and Tree Sprayer may be used. GRAPES: Spray 6 gallons of diluted spray on 1,000 sq. ft.

DILUTIONS: 6 level tablespoons per gallon of water. One pound per 10 gallons of water.

DUSTING—This material is ready for use when applied by a duster. Apply the material directly on the tree. Cover the fruit and foliage thoroughly with a thin coat of dust and apply when air is still. For timing of applications follow directions for spraying.

SUGGESTED FRUIT SPRAY SCHEDULES								
Time to Spray	Delayed Dormant from first green on bud to 1/2 in. leaves	Pre-Bloom from first pink to first bloom	Petal-Fall when last of the petals have fallen	Shuck Fall when half of the shucks have fallen from fruit	Cover Sprays 2 to 4 sprays, depending on local conditions apply at 2 week intervals		Pre-Harvest 2 to 3 weeks before harvest	Post Harvest Immediately after harvest
					Begin 10 days after petal fall	Begin 10 days after shuck fall		
APPLE	X	X	X		X			
PEAR		X	X		X			
CHERRY		X	X	X		X		X
							X	

X indicates when to spray.

CAUTION: Keep out of reach of children. May be harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eyes get medical attention. Wash well before eating or smoking. Do not apply or allow to drift into areas occupied by unprotected persons or animals. If spilled, flush area with plenty of water. Do not store diluted material, make new dilution for each use. Dispose of this container when empty.

Do not apply within 10 days before or after an oil spray treatment. Do not apply to Pears, Grapes, Strawberries and Blueberries closer than 1 day to harvest., Apples, Cherries, Plums and Prunes closer than 3 days to harvest and Peaches closer than 7 days to harvest. Do not add Sulphur to the spray mixture when applying this product to Red Delicious and Baldwin Apples, as the combination of Sulphur and Copper may cause spotting.

Time to Spray	from first green on bud to 1/2 in. leaves	from first pink to first bloom	when last of the petals have fallen	the shucks have fallen from fruit	Intervals		2 to 3 weeks before harvest	Immediately after harvest
					Begin 10 days after petal fall	Begin 10 days after shuck fall		
APPLE	X	X	X		X			
PEAR		X	X		X			
CHERRY		X	X	X		X		X
							X	

X Indicates when to spray.

CAUTION: Keep out of reach of children. May be harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eyes get medical attention. Wash well before eating or smoking. Do not apply or allow to drift into areas occupied by unprotected persons or animals. If spilled, flush area with plenty of water. Do not store diluted material, make new dilution for each use. Dispose of this container when empty.

Do not apply within 10 days before or after an oil spray treatment. Do not apply to Pears, Grapes, Strawberries and Blueberries closer than 1 day to harvest., Apples, Cherries, Plums and Prunes closer than 3 days to harvest and Peaches closer than 7 days to harvest. Do not add Sulphur to the spray mixture when applying this product to Red Delicious and Baldwin Apples, as the combination of Sulphur and Captan may cause spotting, yellowing and dropping of leaves. Injury may occur on Bosc Pears in the Northeast under certain conditions.

This product is toxic to fish. Do not apply near ornamental fish ponds or where runoff is likely to occur.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

Carbaryl

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl methylcarbamate)	4.0%
Carbaryl (1-naphthyl methylcarbamate)	7.5%
Carbaryl (1-naphthyl methylcarbamate)	3.0%
Carbaryl (1-naphthyl methylcarbamate)	85.5%

*TM of Union Carbide Corp. for 1-naphthyl methylcarbamate.

Sold by:

NET WEIGHT ONE POUND

USA Reg. No. 5535-7

Nov 8 1967

REAR PANEL

GRO-WELL FRUIT TREE SPRAY is a combination insecticide and fungicide for the control of insects and diseases on APPLE, PEACH, CHERRY, PEAR and certain other fruit trees.

GRO-WELL FRUIT TREE SPRAY may be used to control: Green Apple Aphids, Pear Psylla Aphid, Codling Moth, Apple Maggot, European Red Mites, Clover Mites, Two Spotted Mites, Williamette Mites, Eye Spotted Bud Mite, Pear Leaf Roller, Apple Red Bug, Pear Psylla, Pear Midge, Pear Plant Bug, Tarnished Plant Bug; on APPLES and PEARS: Oriental Fruit Moth, Pear Curculio, Peach Tree Borer, Cottony Peach Scale, Japanese Beetle on PEACHES, European Red Mite on PLUMS and PRUNES, Black Cherry Aphid, Two Spotted Mites and Tent Caterpillar on CHERRIES; Apple Scab, Brown Rot, Leaf Spot on APPLES, PEACHES, PLUMS, PRUNES and CHERRIES; Cherry Fruit Fly, Cherry Fruitworm and Mealy Plum Aard on CHERRIES, PLUMS and PRUNES; Grape Leafhoppers, Grape Leaf Skeletonizer, Grape Berry Moth, Japanese Beetles, Red Banded Leaf Miner and Downy Mildew on Grapes; Spittlebug, Leaf Roller and Strawberry Weevil on Strawberries; Blueberry Maggot, Cherry Fruitworm and Japanese Beetles on Blueberries.

NOTE: For Peach Tree Borer, make two sprays starting about July 7-15 and another about August 1-10. Consult Extension Authorities for exact timing in your area. Where disease control close to harvest is necessary, apply a suitable fungicide alone in a manner which will not leave residues in excess of established tolerances. Where mite control is a serious problem add 1 tablespoonful of GRO-WELL KELTHANE per gallon of spray mixture.

DIRECTIONS: SPRAYING—Add the required amount of material to water and stir thoroughly. Pour the mixture into the sprayer and agitate while spraying in order to obtain a thorough and even coverage. In a small sprayer, shake often for agitation. A sprayer that attaches to the hose such as the Gro-Well Shrub and Tree Sprayer may be used. GRAPES: Spray 6 gallons of diluted spray on 1,000 sq. ft.

DILUTIONS: 6 level tablespoons per gallon of water. One pound per 10 gallons of water.

DUSTING—This material is ready for use when applied by a duster. Apply the material directly on the tree. Cover the fruit and foliage thoroughly with a thin coat of dust and apply when air is still. For timing of applications follow directions for spraying.

SUGGESTED FRUIT SPRAY SCHEDULES								
Time to Spray	Delayed Dormant from first green on bud to 1/2 in. leaves	Pre-Bloom from first pink to first bloom	Petal-Fall when last of the petals have fallen	Shuck Fall when half of the shucks have fallen from fruit	Cover Sprays 2 to 4 sprays, depending on local conditions apply at 2 week intervals		Pre-Harvest 2 to 3 weeks before harvest	Post Harvest Immediately after harvest
					Begin 10 days after petal fall	Begin 10 days after shuck fall		
APPLE	X	X	X		X			
PEAR		X	X		X			
CHERRY		X	X	X		X		X

Indicates when to spray.

CAUTION: Keep out of reach of children. May be harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eyes get medical attention. Wash well before eating or smoking. Do not apply or allow to drift into areas occupied by unprotected persons or animals. If spilled, flush area with plenty of water. Do not store diluted material, make new dilution for each use. Dispose of this container when empty.

Do not apply within 10 days before or after an oil spray treatment. Do not apply to Pears, Grapes, Strawberries and Blueberries closer than 1 day to harvest., Apples, Cherries, Plums and Prunes closer than 3 days to harvest and Peaches closer than 7 days to harvest. Do not add Sulphur to the spray mixture when applying this product to Red Delicious and Baldwin Apples, as the combination of Sulphur and Captain may cause spotting, yellowing and dropping of leaves. Injury may occur on Bose Pears in the Northeast under certain conditions.

This product is toxic to fish. Do not apply near ornamental fish ponds or where runoff is likely to occur.

PEAR		X	X		X		
CHERRY		X	X	X		X	X
			X	X		X	X

X indicates when to spray.

CAUTION: Keep out of reach of children. May be harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water; for eyes get medical attention. Wash well before eating or smoking. Do not apply or allow to drift into areas occupied by unprotected persons or animals. If spilled, flush area with plenty of water. Do not store diluted material, make new dilution for each use. Dispose of this container when empty.

Do not apply within 10 days before or after an oil spray treatment. Do not apply to Pears, Grapes, Strawberries and Blueberries closer than 1 day to harvest., Apples, Cherries, Plums and Prunes closer than 3 days to harvest and Peaches closer than 7 days to harvest. Do not add Sulphur to the spray mixture when applying this product to Red Delicious and Baldwin Apples, as the combination of Sulphur and Captan may cause spotting, yellowing and dropping of leaves. Injury may occur on Bosc Pears in the Northeast under certain conditions.

This product is toxic to fish. Do not apply near ornamental fish ponds or where runoff is likely to occur.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS

Carbaryl	(1-naphthyl N-methylcarbamate)	4.0%
	(1-naphthyl N-methylcarbamate)	7.5%
		3.0%
		85.5%

* 0.0% active ingredient as calcium salt of diethyl malathionate
 * TM of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

Sold by: _____

NET WEIGHT 4 1/2 POUNDS USDA Reg. No. 55357

nov 8 1967 5535-7

**PEAR
PANEL**

GRO-WELL FRUIT TREE SPRAY is a combination Insecticide and Fungicide for the control of Insects and Diseases on APPLE, PEACH, CHERRY, PEAR and certain other fruit trees.

GRO-WELL FRUIT TREE SPRAY may be used to control: Green Apple Aphids, Woolly Apple Aphid, Codling Moth, Apple Maggot, European Red Mites, Clover Mites, Two Spotted Mites, Willamette Mites, Eye Spotted Bud Moth, Fruit Tree Leaf Roller, Apple Red Bug, Pear Psylla, Pear Midge, Pear Plant Bug, Tarnished Plant Bug on APPLES and PEARS; Oriental Fruit Moth, Plum Curculio, Peach Tree Borer, Cottony Peach Scale, Japanese Beetle on PEACHES, European Red Mite on PLUMS and PRUNES, Black Cherry Aphid, Two Spotted Mites and Tent Caterpillar on CHERRIES; Apple Scab, Brown Rot, Leaf Spot on APPLES, PEACHES, PLUMS, PRUNES and CHERRIES; Cherry Fruit Fly, Cherry Fruitworm and Mealy Plum Aphid on CHERRIES, PLUMS and PRUNES. Grape Leathoppers, Grape Leaf Skeletonizer, Grape Berry Moth, Japanese Beetles, Red Banded Leaf Roller and Downey Mildew on Grapes. Spittlebug Leaf Roller and Strawberry Weevil on Strawberries. Blueberry Maggot, Cherry Fruitworm and Japanese Beetles on Blueberries

NOTE: For Peach Tree Borer, make two sprays starting about July 7-15 and another about August 1-10. Consult Extension Authorities for exact timing in your area. Where disease control close to harvest is necessary, apply a suitable fungicide alone in a manner which will not leave residues in excess of established tolerances. Where mite control is a serious problem add 1 tablespoonful of GRO-WELL KELTHANE per gallon of spray mixture.

DIRECTIONS: SPRAYING—Add the required amount of material to water and stir thoroughly. Pour the mixture into the sprayer and agitate while spraying in order to obtain a thorough and even coverage. In a small sprayer, shake often for agitation. A sprayer that attaches to the hose such as the Gro-Well Shrub and Tree Sprayer may be used. GRAPES: Spray 6 gallons of diluted spray on 1,000 sq. ft.

DILUTIONS: 6 level tablespoons per gallon of water. One pound per 10 gallons of water.

DUSTING—This material is ready for use when applied by a duster. Apply the material directly on the tree. Cover the fruit and foliage thoroughly with a thin coat of dust and apply when air is still. For timing of applications follow directions for spraying.

SUGGESTED FRUIT SPRAY SCHEDULES								
Time to Spray	Delayed Dormant from first green on bud to 1/2 in. leaves	Pre-Bloom from first pink to first bloom	Petal-Fall when last of the petals have fallen	Shuck Fall when half of the shucks have fallen from fruit	Cover Sprays 2 to 4 sprays, depending on local conditions apply at 2 week intervals		Pre-Harvest 2 to 3 weeks before harvest	Post Harvest Immediately after harvest
					Begin 10 days after petal fall	Begin 10 days after shuck fall		
APPLE	X	X	X		X			
PEAR		X	X		X			
CHERRY		X	X	X		X		X
PEACH PLUM	X	X	X	X		X	X	

apply or allow to drift into areas occupied by unprotected persons or animals. If spilled, flush area with plenty of water. Do not store diluted material, make new dilution for each use. Dispose of this container when empty.

Do not apply within 10 days before or after an oil spray treatment. Do not apply to Pears, Grapes, Strawberries and Blueberries closer than 1 day to harvest., Apples, Cherries, Plums and Prunes closer than 3 days to harvest and Peaches closer than 7 days to harvest. Do not add Sulphur to the spray mixture when applying this product to Red Delicious and Baldwin Apples, as the combination of Sulphur and Captan may cause spotting, yellowing and dropping of leaves. Injury may occur on Bosc Pears in the Northeast under certain conditions.

This product is toxic to fish. Do not apply near ornamental fish ponds or where runoff is likely to occur.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the ~~buyer assumes all~~ risks if directions given herewith are not strictly adhered to.

ACCEPTED
 nov 8 1967
 5535-7
 4.0%
 7.5%
 3.0%
 85.5%
 NET WEIGHT ONE POUND
 USDA Reg. No. 5535-7

ACTIVE INGREDIENTS:

**Malathion

Captan (N-trichloromethylthio tetrahydrophthalimide)
 (1-Naphthyl N-Methylcarbamate)

INERT INGREDIENTS:

**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

*TM of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate

Carbaryl

Sold By:

Union Carbide Corp., 270 Park Ave., New York 17, N.Y.