

GRO-WELL ROSE & GARDEN FUNGICIDE

GRO-WELL ROSE & GARDEN FUNGICIDE will control certain diseases on Roses, Fruit Trees, Flowers and Vegetables. May be mixed with GRO-WELL Liquid Insecticide to control both insects and diseases.

Add 3 level tablespoonfuls to each gallon of water (8 oz. per 10 gallons of water). Stir thoroughly. Stir or agitate while spraying in order to obtain uniform coverage. Spray plants thoroughly being sure to cover underside of leaves. A sprayer which attaches to the hose, such as the Gro-well Sprayer, may be used.

ROSES AND FLOWERS: Will control Black Spot, Powdery Mildew, Clover and Red Spider Mites on ROSES, Rust and Leaf Spot on CARNATIONS; Flower Blight on CHRYSANTHEMUMS; Powder Mildew on DELPHINIUM, DAHLIAS, HYDRANGEAS and SNAPDRAGONS. Apply as above at first sign of disease and repeat at 7 to 10 day intervals and after heavy rains.

FRUIT TREES: Will control Scab, Powdery Mildew, Frogeye, Brooksfuit Spot, Sooty Blotch, Fly Speck, Black Rot and Bitter Rot on APPLES; Scab on PEARS; Brown Rot and Scab on PEACHES; Downy Mildew and Black Rot on GRAPES. Note: Begin application at Pre-Bloom and continue at 7 to 10 day intervals making last application 21 days before harvest on APPLES, PEARS, GRAPES; 45 days on PEACHES.

VEGETABLES: Will control Powdery and Downey Mildew, Anthracnose and Angular Leaf Spot on CANTELOUPES, CUCUMBERS AND SQUASH. Use 4 oz. of Garden Fungicide per 5 gallons of water to cover 1000 Ft. of garden area. Begin application at first signs of disease and repeat at 5 to 7 day intervals or as needed to maintain control. Note: Make last application 7 days before harvest.

CAUTION: Keep out of reach of children and domestic animals. Avoid inhalation of Dust or Spray mist; avoid contact with skin. May be a skin irritant and sensitizer. Wash well after using. Avoid contamination of Feed and Foodstuffs. Do not add Sulphur to the spray mixture when applying to Delicious and Baldwin Apples as the combination may cause Spotting, Yellowing and dropping of leaves. Injury may occur on Bosc Pears in the Northeast under certain conditions. Where disease control close to harvest is necessary apply a suitable Fungicide alone in a manner which will not leave residue in excess of established tolerances.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS:

Captan (N-Trichloromethylmercapto-4-Cyclohexene-1-2-Dicarboximide)	20.00%
*Dinitro (1-Methylheptyl) phenyl crotonate	1.80%
Other Nitrophenols and derivatives chiefly	
Dinitro (1-Methylheptyl) phenol20%
INERT INGREDIENTS	78.00%
*Karathane—Reg. TM Rohm & Haas Co.	100.00%


Sold By: NET WEIGHT 8 OUNCES U.S.D.A. REG. NO. 5535-33
 J. & L. ADIKES, INC., JAMAICA, NEW YORK

Narvon Clay (Narvon Mines)

TOTAL INERTS



CAPTAN & KARATHANE*
 For control of certain diseases on
ROSES, FLOWERS, FRUIT TREES & VEGETABLES



CAUTION:
 KEEP OUT OF REACH OF CHILDREN
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS

2.0%
 49.0%

 78.0%

ACCEPTED
 JUL 9 1968
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 5535-33 SUBJECT
 TO ATTACHED COMMENTS