

EPA Reg. No. 5535-33

JAMAICA, N.Y. 11423

STORAGE. Store in ongoal can in a locked storage area, every from other pet-bories. Instituter, lood, and lead. Keep can bythy closed after opening it broken or

#R.g. 5535-33

GRO-WELL* ROSE & GARDEN FUNGICIDE will control certain diseases on ROSES_FLOWERS_FRUIT TREES and VEG-

Mix the required dosage with water (as noted), stirring thoroughly and agitating while spraying in order to obtain a more uniform coverage. Spray plants thoroughly, being sure to cover underside of leaves. A sprayer which attaches to a hose, such as the GRO-WELL* SHRU8 & TREE SPRAYER, may be used.

CAUTION: Do not add Sulphur to the spray mixture when applying to Delicious and Baldwin Apples, as the combination may cause spotting, yellowing and dropping of leaves. Injury might occur on Bosc Pears in the Northeast under certain conditions. Where disease control close to harvest is necessary, apply a suitable fundicide alone in a manner which will not leave residues in excess of established tolerances.

WARRANTY STATEMENT

51610-2M-10-B5

Follow directions carefully, as use, storage and handling of this material is beyond the control of the selier. The buyer assumes at nsks & directions given herewith are not strictly achered to. Seller gives no other or further Warranty, expressed or implied as to the merchantability or any other aspect of

NOTES: ① Apply at first sign of dis-ease. Repeat at 7 to 10 day intervals and after heavy rains. (2) Begin application at pre-bloom and continue at 7 to 10 day intervals. (1) Do not apply within 21 days of harvest. (2) Do not apply within 45 days of harvest. (§) Apply diluted spray to an area of 1,000 sq. ft. (20' x 50') at first sign of disease. Receat at 5 to 7 day intervals or as needed to maintain control. Do not apply within 7 days of harvest,

1 As Listed