



CAUTION
 KEEP OUT OF REACH OF CHILDREN
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:
 Thiram (Tetramethylthiuram Disulfide) 6.20%
 Cadmium Chloride* 1.20%
INERT INGREDIENTS: 92.60%
 *Total Cadmium 0.69%

SOLD BY:
J. & L. ADIKES, INC. JAMAICA, N.Y.

U. S. D. A. Reg. No. 5535-34

CCER
JUN - 35 - 1968
5535-34

GRO-WELL

LAWN FUNGICIDE

This easy to apply dry form is for the prevention and control of Dollar Spot, Copper Spot, Leaf Spot (Helminthosporium) Curvularia, Brown Patch and Snow Mold, on lawns containing Kentucky Blue, including Merion and other strains, the various Fescues, Bents and most other grasses. Contains Thiram and Cadmium Chloride, plus two additional ingredients which improve the prevention and control of diseases.

SPECIAL INSTRUCTIONS:

After applying Gro-Well Lawn Fungicide, do not water or mow treated lawn area for 24 hours. Application may be made at any time. The ideal time would be early morning when dew is on the grass, causing material to adhere to grass blades, thus improving results. If desired, area may be watered just prior to application and this will accomplish the same result. Gro-Well Lawn Fungicide may be used as a preventative against lawn diseases by applying this bag to 5000 square feet. It may also be used to protect lawns against heavy damage caused by existing diseases, by reducing the further spread of the disease. Use at heavy application rate for this purpose.

EARLY FALL APPLICATIONS:

With the coming of cooler nights, Dollar Spot and Copper Spot diseases may appear again. Prevent infestation with treatments at 10 day to 2 week intervals.

WINTER APPLICATION:

Snow Mold: Apply at the heavy application rate in the late fall, before the first snowfall and repeat again at the same rate in January.

WARNING:

Avoid contact with eyes, skin or clothing. In case of contact, wash exposed areas thoroughly with water to avoid

cide may be used as a preventative against lawn diseases by applying this bag to 5000 square feet. It may also be used to protect lawns against heavy damage caused by existing diseases, by reducing the further spread of the disease. Use at heavy application rate for this purpose, applying this bag to 2500 square feet of lawn area.

SPRING PROGRAM:

Leaf Spot, usually found in early Spring or Dollar Spot, usually prevalent in late Spring, can cause grass plants to die. To prevent this damage, apply Lawn Fungicide soon after second mowing in early Spring at 10 day to 2 week intervals. Three to four applications are usually necessary. To arrest these diseases if they are already attacking lawns, use heavy application rate. Repeat as necessary.

SUMMER APPLICATION:

Prevention of Brown Patch or Curvularia diseases, which become active when nights are warm and humidity high, may be obtained by applying Lawn Fungicide at the normal rate at 10 day to 2 week intervals through the hot, humid months. If diseases are present and damage is visible use heavy application rate. Repeat as necessary.

Where heavy application rate is needed, go over area twice at above settings.

NOTE: Above spreader settings are approximate. Allowances must be made for condition of spreader, humidity, speed at which applied, and roughness of area treated.

NET WEIGHT 8 LBS.

first snowfall and repeat again at the same rate in January.

WARNING:

Avoid contact with eyes, skin or clothing. In case of contact, wash exposed areas thoroughly with water to avoid irritation. Avoid breathing dust, since inhalation may prove harmful. Keep out of reach of children. **INTERNAL:** If accidentally swallowed, give raw white of egg. Induce vomiting with mixture of salt, soap or mustard in warm water. Call physician at once.

NOTE:

We make no warranty of any kind expressed or implied concerning the suitability or results of this product. Buyer, user or handler assumes all risks, whether or not directions are followed.

SPREADER SETTINGS:

CYCLONE	2 1/2
SCOTTS	5 1/2
LAWN BEAUTY	4 1/2
OTHERS	3/16"