


PROD. # 51250

GRO-WELL.

FERBAM

A FUNGICIDE

CONTROLS BLACK SPOT on ROSES,
 BLIGHTS, RUST AND OTHER DISEASES
 on FLOWERS, SHRUBS, TREES
 AND CERTAIN VEGETABLES*



MAKES UP TO
 75 GALLONS OF SPRAY

ACTIVE INGREDIENT:

Ferbam (Ferric Dimethyl Dithiocarbamate)..... 76.00%

INERT INGREDIENTS:

24.00%

TOTAL 100.00%

ACCEPTED

OCT 27 1983

U.S. Dept. of Agriculture
 Federal Insecticide, Fungicide, and Rodenticide Act
 as amended for the pesticide
 registration act of 1972
 EPA Reg. No. 263-100-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY INFORMATION

NET WEIGHT
 8 oz (1/2 lb)

*As listed

SINCE 1855 SOLD BY: J. & L. ADIKES, INC.
 JAMAICA, N.Y. 11423

REAR PANEL

FERBAM

TYPE SIZE

----- PRECAUTIONARY STATEMENTS

----- HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: KEEP OUT OF REACH OF CHILDREN. May cause irritation of eyes, nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. In case of contact, flush with plenty of water; for eyes, contact physician. Keep away from fire or sparks. Store away from domestic animals, feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using. Use strictly in accordance with label cautions and directions. Apply this product only as specified on this label.

----- STORAGE AND DISPOSAL

STORAGE: Store in original can in a locked storage area, away from domestic animals, other pesticides, fertilizer, food, and feed. Keep can tightly closed after opening. If broken or spilled, sweep up and discard in trash.

PESTICIDE DISPOSAL: Securely wrap original can in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty can. Rinse thoroughly before discarding in trash.

----- WARRANTY STATEMENT

Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given herewith are not strictly adhered to. Seller gives no other or further warranty, expressed or implied as to the merchantability or any other aspect of products sold.

BEST DOCUMENT AVAILABLE

GRO-WELL® FERBAM is a fungicide which will control many diseases on a large variety of plants including TREES, SHRUBS, FLOWERS, FRUITS and VEGETABLES. See list for specific diseases controlled. A sprayer which attaches to a hose, such as the GRO-WELL® SHRUB & TREE SPRAYER, may be used. Follow mixing directions on sprayer. Cover plants thoroughly with spray, being sure to spray the underside of leaves as well.

DOSAGE LISTED TO BE MIXED WITH 1 GALLON OF WATER

FOR USE ON	SEE NOTE	TO CONTROL	DOSAGE (Application Rate)
FLOWERS-SHRUBS-TREES: Asters, Begonias, Gardenia, Geraniums, Hydrangea, Ivy, Laurel	8	Leaf Spot	2 level tablespoonsful
Azaleas		Leaf Scorch	
Camellias	1-8	Petal Blight	
Camations		Alternaria Blight, Rust, Septoria Leaf Spot	
Chrysanthemums		Rust, Septoria Leaf Spot	
Cyclamen	8	Botrytis Rot	
Junipers		Cedar Rot	
Roses		Black Spot, Rust	
Snagdragons		Leaf Spot, Rust	
Tulips		Botrytis Blight	
FRUITS:			
Apples	2-5-9	Apple Blotch, Bitter Rot, Black Rot, Blossom End Rot, Brooks Spot, Cedar Apple Rust, Fly Speck, Sooty Blotch	
Apricots	3-10	Brown Rot Blossom Blight, Green Rot, Shothole	
Blueberries	4-11	Botrytis Blight	
Grapes	4-5-12	Black Rot	
Blackberries, Raspberries	4-13	Anthracoese, Rust, Spur Blight	
Peaches	3-14	Brown Rot Blossom Blight, Leaf Curl, Scab, Shothole	
Pears	5-15	Fabraea Leaf Blight, Fruit Spot, Scab, Shothole	
Plums, Prunes	5-16	Brown Rot	
Cherries	5-17	Brown Rot, Leaf Spot	
VEGETABLES:			
Cabbage	6-18	Downy Mildew(Plant bed treatments only)	
Lettuce	6-19	Damping Off(Plant bed treatments only)	
Celery	8	Leaf Spot	
Snapbeans, Stringbeans	7-8	Anthracoese	

NOTES: ① Apply also to the soil and litter around plants while they are in bloom. ② Do not use on certain tender varieties such as Golden Delicious as it may contribute to fruit russeting. ③ Do not apply within 21 days of harvest. ④ Do not apply closer than 40 days of harvest. ⑤ Do not apply closer than 7 days of harvest. ⑥ Remove residue by stripping, trimming and washing. ⑦ Do not apply within 4 days of harvest and do not feed treated plants to livestock. ⑧ Use before diseases are apparent and spray every 7-14 days as long as there is a disease problem. ⑨ To control Rust, apply in pre-bloom, calyx & early cover sprays; for Black Rot, apply at petal fall & repeat in cover sprays; for other listed diseases, apply in cover sprays. ⑩ To control Brown Rot, apply before, during & after bloom and in cover sprays as necessary; for Green Rot, apply at full bloom & petal fall; for Shothole, apply after harvest, repeat in Spring at end of swell & shuck shed. ⑪ Apply just before blooming, repeat at full bloom & 1-3 weeks later. ⑫ Apply just before blooms, just after bloom & 2 weeks before berries touch. ⑬ Apply at delayed dormant stage, just before bloom & at petal fall. ⑭ For Brown Rot & Scab, apply before, during & after bloom and repeat as necessary; for Shothole, apply after harvest but before leaves fall, repeat in Spring at popcorn & shuck shed stages; for Leaf Curl, in Western States, apply in Fall after leaves drop & in early Spring dormant period; for Eastern States, make one dormant season application. ⑮ Apply in pink & calyx stages & through second cover sprays. ⑯ Apply in popcorn, bloom, petal fall & first and second cover sprays. ⑰ For Brown Rot, apply at pink bud stage, repeat at full bloom, petal fall & 1 or more cover sprays as necessary; for Leaf Spot, make 2-3 applications before harvest & one application after harvest. ⑱ Apply when plants emerge & repeat 2-3 times per week for 3-4 applications. ⑲ Apply when plants emerge & repeat at 7-10 day intervals. GENERAL: Contact State Extension Service or Experiment Station experts for further information on timing of applications.