

GRO-WE L CRAB-N-WELD is a dry granular material which when applied to your lawn with a spreader will kill CRABGRASS and certain broadleaf weeds, such as PLANTAIN, DANDLION, PURSLANE, YARROW, COMMON CHICKWEED and GROUND IVY. May be applied at any time weeds are prevalent, as long as cautions given below are followed. For Crabgrass two to three applications, 10 to 14 days apart, may be necessary for best control. For broadleaf weeds one or two applications may be sufficient. Since crabgrass continues to germinate throughout the summer months, repeated treatments may be necessary to kill new growth.

To eliminate the possibility of future crabgrass infestations, it is recommended that an application of GRO-WE L PRE-VENT[®], a pre-emergent crabgrass control be applied each year to prevent the germination of crabgrass seedlings.

WHEN & HOW TO APPLY:

Thoroughly moisten area to be treated before application so that chemical will adhere to the leaves. This product should remain on the leaf surface for 48 hours. Do not water during this period. After the 48 hours water thoroughly to reduce the danger of injury to desirable grasses. Thereafter, water is required for good growth of turf grasses.

CAUTION: KEEP OUT OF REACH OF CHILDREN & DOMESTIC ANIMALS. HARMFUL IF SWALLOWED. Keep children and pets off treated area until completely dry. Avoid inhalation of dust and contact with skin, eyes and clothing. Wash thoroughly with soap and warm water after using and before smoking or eating. In case of contact with eyes, flush with water profusely and contact doctor. If taken internally, induce vomiting and contact doctor. If spilled, dispose of immediately. Do not store near other lawn or garden supplies. Do not contaminate feed or foodstuffs. Do not ~~use~~ re-use container, dispose of in a safe manner when empty, either by burying with other waste or in a trash can which can be tightly sealed.

CAUTION: Do not use on newly seeded lawns until after third mowing. For best results do not re-seed until 3 weeks after last application. May cause damage to Bent, Dichondra, St. Augustine grass, Centipede grass and Clover lawns. If weed killing is desired in hot weather (over 90°) thoroughly soak lawn with water and apply at half rate, otherwise injury might occur. Repeat treatment in 2 weeks if necessary. Apply when air is still to avoid getting material into flower or vegetable beds. Vapors from this product may cause injury to susceptible plants in the immediate vicinity.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

SPREADER SETTINGS

Cyclone..... 2-1/2
Scotts..... 3-1/2
Lawn Beauty..... 5

NOTE: Above spreader settings are approximate. Allowances must be made for condition of spreader, humidity, speed at which applied and roughness of area treated.

FRONT PANEL

Inc. 1955

G R O W E L L

W E E D - N - C H I C K

Apply by spreader
COVERS 5000 sq. ft.

KILLS CHICKWEED, DANDELION,
PLANTAIN & OTHER LAWN WEEDS.

Active Ingredients:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*.....	2.45%
*2,4-D Acetic Acid equivalent 2.00%	
Butoxypolypropoxypropyl ester of Silvex**.....	1.85%
**Silvex (2-(2,4,5-Trichlorophenoxy)propionic acid)	
equivalent 1.30%	

Inert Ingredient..... 95.70%

TOTAL 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

SOLD BY:

J. J. ABLETT, INC.

JAMAICA, NEW YORK 11426

U.S.P.A. REG. NO. 5535-41