

REAR PANEL

GRO-WELL WEED-N-CHICK is a dry granular material which when applied to your lawn will kill most broadleaf lawn weeds such as Dandelion, Plantain, Chickweed (common & mouse-ear), Knotweed, Purslane, Yarrow, Ground Ivy, Pennywort and many others. For broadleaf weeds one application may be sufficient, however, if weeds are not completely killed repeat application in three weeks.

WHEN & HOW TO APPLY:

May be used at any time as long as cautions are followed. Best results are obtained when soil is moist and lawn is in a good growing condition. Thoroughly moisten area to be treated before application so that chemical will adhere to the leaves. WEED-N-CHICK should remain on the weeds for 48 hours so do not water until 48 hours after application; then water thoroughly to reduce the danger of injury to desirable grasses.

CAUTION:

Do not use on newly seeded lawns until third mowing. For best results do not re-seed until 2 weeks after last application. Avoid applying during periods of drought as temporary browning of desirable grasses may occur. If weed killing is desired in hot weather (over 90°) or on a clover or bent lawn thoroughly soak lawn with water and apply at half rate, otherwise damage may result. Do not use on Dichondra and St. Augustine grass as injury might occur. Repeat treatment in two weeks if necessary. Apply when air is still to avoid getting material into flower or vegetable beds or under the spread of trees or shrubs as vapors from this product may cause injury to susceptible plants in the immediate vicinity. Do not store near other lawn or garden supplies.

WARNING: Keep out of reach of children and domestic animals. Keep children and pets off until treated area is dry. Avoid inhalation of dust and contact with skin, eyes or clothing; wash off with soap and warm water. In case of contact with eyes, flush with water profusely and contact doctor. If taken internally, induce vomiting and contact doctor. If spilled, dispose of immediately. Dispose of this container when empty.

NOTICE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

SPREADER SETTINGS:


- Cyclone.....2-1/2
- Scotts.....1-1/2
- Lawn Beauty.....1

NOTE: Above spreader settings are approximate. Allowances must be made for condition of spreader, humidity, speed at which applied, and roughness of area treated.

NET WEIGHT

STATEMENT OF INERT INGREDIENTS

Vermiculite	10.50%
Water	5.20%
	15.70%



GRO-WELL

ACCEPTED
7-26-68
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR SALE IN THE UNITED STATES
EPA REG. NO. 5535-46

Comments

FEED-N-WEED

GRO-WELL FEED-N-WEED will kill most broad-leaved weeds such as Dandelion, Plantain, Chickweed (common and mouse-ear), Pennywort, Ground Ivy plus certain others.

The 10-6-4 fertilizer contained in Feed-N-Weed will help to fill in areas where weeds have been killed as well as improve the general appearance of your lawn.

WHEN & HOW TO APPLY:

Feed-N-Weed should be applied in late Spring, Summer or Fall.

FOR BEST RESULTS:

1. Lawn should be cut a day or two before application.
2. Soil should be thoroughly moistened from either rainfall or heavy watering 24 hours before treatment, so that grasses are in good growing condition.
3. Weeds should be treated immediately before application so that material will adhere to the weeds.
4. Do not water for 48 hours after application, then water thoroughly to reduce the danger of injury to desirable grasses in the event an excessive amount has been applied to any area.
5. Do not mow lawn for 48 hours after application.

grasses may occur in periods of drought, follow Step Two closely to reduce this danger if weed killing is desired during these periods.

WARNING:

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not spread or allow to drift onto desirable shrubs, flowers, trees and vegetables or into lakes, streams or ponds. Keep out of reach of children and pets. Avoid inhalation of particles and contact with skin, eyes or clothing. Wash well with soap and water after using. If accidentally taken internally, induce vomiting, contact doctor immediately. If gotten into eyes flush profusely with water and contact doctor. Do not store near seeds, fertilizers, insecticides or fungicides.

7-26-68

5335-46

CAUTION:

Areas may be seeded three weeks after application. Do not apply to newly seeded lawns until after the second or third mowing and grasses have become established. Feed-N-Weed may cause damage to Bent and Clover lawns. Damage or temporary browning to desirable

NOTICE:

We make no warranty of any kind expressed or implied concerning the suitability or results of this product. Buyer, user or handler assumes all risks whether or not directions are followed.

SPREADER SETTINGS

SCOTTS	6 +
LAWN BEAUTY	5 3/4
CYCLONE	3
OTHERS	3/16"

NOTICE: Above spreader settings are approximate. Allowances must be made for condition of spreader, humidity, speed at which applied and roughness of area to be treated.


2-26-68
5530-46

NET WEIGHT 20 LBS.

Kills



BROAD-LEAVED
PLANTAIN



DANDELION



MOUSE-EAR
& COMMON
CHICKWEED

AND
MANY OTHER
ANNOYING
LAWN WEEDS

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ANALYSIS - FERTILIZER

Nitrogen (N)	10.00 %
Available Phosphoric Acid (P ₂ O ₅)	6.00 %
Potash (Soluble in water) (K ₂ O)	4.00 %
Potential Acidity equivalent to 600 pounds Calcium Carbonate per ton.	

7-24-47

5535-46

INGREDIENTS - HERBICIDE

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D, Dichlorophenoxy-Acetic Acid*	1.00 %
2,4-D Acid Equivalent .84 %	
Solihexacyl ester of Silvex (2-(2,4,5-trichlorophenoxy) propionic acid)**	0.55 %
**2-(2,4,5-trichlorophenoxy) propionic acid equiv. 0.37 %	
INERT INGREDIENTS (Herbicide Content)	98.45 %

SOLD BY: **J. & L. ADIKES, INC. JAMAICA, N.Y.** U.S. D. A. Reg. No. 5636-46