


FRONT PANEL

[182]



INSECTICIDE • FUNGICIDE • MITICIDE
FOR THE CONTROL OF MANY INSECTS AND DISEASES

ACCEPTED

REAR PANEL

[282]

MAKES 10 GALLONS OF SPRAY

GRO-WELL MULTI-PURPOSE DUST or SPRAY is effective for control of the following **INSECTS** and **DISEASES** on **ROSES, FRUIT TREES, SHRUBS, TREES** and **VEGETABLES**.

DIRECTIONS

DUSTING: Apply to Roses, Fruit Trees, Shrubs and Vegetables before insects become numerous and before diseases become apparent. Protect plants through the growing season by dusting at 7 to 14 day intervals and after heavy rains. Cover foliage thoroughly, but do not use in excess. Pay particular attention to cover underside of leaves. On Fruit Trees, apply the material directly to the tree when the air is still. For timing of applications, follow Fruit tree spray and dust schedule.

SPRAYING: Add the required amount of material to water and stir thoroughly. Pour the mixture into the sprayer and agitate while spraying in order to obtain a thorough and even coverage. In a small sprayer, shake often for agitation.

DILUTIONS: 6-7 level tablespoons per gallon of water. One pound per 10 gallons of water, 10-12 pounds per 100 gallons of water.

ROSES

Aphids, Spider Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Flea Beetles, Thrips, Tarnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on **ROSES, GLADIOLUS, DAHLIAS, SWEET PEAS, ASTER** and **CHRYSANTHEMUMS**. Black Spot, Powdery Mildew, Rust, Leaf Spot and Botrytis Flower Blight of **ROSES, TUBEROUS BEGONIAS, CARNATIONS, CHRYSANTHEMUMS** and **PEONIES**.

FRUIT TREES

Aphids, Codling Moth, Mites (European Red, Clover, Two Spotted, Willemette), Pear Psylla, Japanese Beetles, Black Cherry Aphid, Tent Caterpillar, Apple Scab, Sooty Blotch, Brooks Spot, Bitter Rot, Fly Speck, Brown Rot, Peach Scab, Leaf Spot on **APPLES, PEARS, PEACHES** and **CHERRIES**.

SHRUBS

Japanese Beetles, Aphids, Spider Mites, Lacebugs, Scale (Azalea, Scurfy, Euonymus, Soft Brown), Bagworms and Boxwood Leaf Miners on **AZALEAS, RHODODENDRONS, ARBORVITAE, JUNIPERS, HYDRANGEAS, EUONYMUS, PIERIS JAPONICA, LILAC** and **BOXWOOD**.

NOTE: For scale insects and Bagworms on **SHRUBS**, make only dusting applications. Do not spray for control of these insects as 2 to 3 times the normal dosage of this product is required per gallon of water. This would result in excessive clogging of spray nozzles. Powdery Mildew on **LILAC** and **EUONYMUS**.

NOTE: Do not use on Boston, Maidenhair, Pteris Ferns as well as species of **Crassula**. Do not apply on **Petunias**.

TREES

Spider Mites, Birch Leaf Miners, Holly Leaf Miners, European Pine Shoot Moth, Tent Caterpillars and Aphids on **PINE, BIRCHES, HOLLY, OAKS, MAGNOLIAS, MAPLES** and **SYCAMORE**.

VEGETABLES

TOMATOES: Fruitworms, Hornworms, Aphids, Spider Mites, Flea Beetles, Colorado Potato Beetles, Early and Late Blight, Anthracnose and Septoria Leaf Spot.

PEPPERS and **EGGPLANTS:** Aphids, Flea Beetle, Colorado Potato Beetles and Lacebugs.

POTATOES: Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers, Early and Late Blight.

PEAS: Pea Aphid.

SUGGESTED FRUIT TREE SPRAY AND DUST SCHEDULE

DELAYED DORMANT (From first green on bud to 1/2 inch leaves on Apple and Peach).

PRE-BLOOM (From first pink to first bloom on Apple, Peach, Pear and Cherry).


PETAL FALL -- (When last petals have fallen) on all fruits listed above.

SHUCK FALL (When half the shucks have fallen from fruit on Cherry and Peach).

COVER SPRAYS (2 to 4 applications) Begin 10 days after petal fall on Apple and Pear. Begin ten days after shuck fall on Cherry and Peach.

CAUTION: Keep out of reach of children and pets. Harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Wash hands thoroughly before eating or smoking. Do not contaminate feed and foodstuffs. Do not apply or allow to drift into areas occupied by unprotected persons or beneficial animals. If spilled, flush area with plenty of water. Dispose of this container when empty. Do not store diluted spray material. Make new dilution for each use.

Do not use Sulphur on apples within 10 days to 2 weeks of an oil spray. Do not make applications when temperature is high as Sulphur may burn foliage. Do not apply to Red Delicious Apples as the combination of Captan and Sulphur may cause foliage injury. Injury may occur on Bosc Pears in the Northeast under certain conditions. Injury may occur to certain varieties of Sweet Cherries, especially in the Northwest. Do not apply to Apples, Cherries, Eggplants and Peas within 3 days of harvest. Do not apply to Pears within 1 day of harvest. Do not apply to Tomatoes within 5 days of harvest. Do not apply to Peaches within 7 days of harvest. After the time limits set above have passed, an effective fungicide should be used alone in a manner which would not leave residues in excess of established tolerances. Do not use on crops to be fed to dairy animals or to animals being finished for slaughter.



SOLD BY:

NET WEIGHT — 1 POUND

U.S.D.A. REG. NO. 55317-2