



For the destruction of weeds and grasses Prior to seeding a new lawn

22.6% Cacodylate (sodium dimethylarsinate) vic Acid Dimethylarsinic Acid) REDIENTS: 3.9% 73.5% ptal Arsenic, all in water soluble form, as elemental arsensic 12.7%. CAUTION REPORT OF REAL HOTE CHILDREN

Covers 3,200 sq. ft.

BREDIENTS:

GRO-WELL REN-O-VATE

Gro-Well Ren-O-Vate is for use in destroying all top growth, both grasses and weeds prior to reseeding for a new lawn, or when used with care, for killing weeds and grasses around shrubs, trees and similar areas where there is unwanted vegetation

Good results will be obtained if applied during the growing season from Spring through Fall on a warm day. Mow lawn at 1 height and rake before chemical is applied. Use 10 ozs. of Ren-O-Vate in 5 gallons of water and apply to 1000 sq. ft of lawn area (20 x 50). A sprayer such as the Gro-Well Shrub & Tree Sprayer, which attaches to a hose may be used. If this type sprayer is used, add additional water to the chemical in the jar so that the jar is filled up to the five gallon mark. This will give you the required 5 gallons of diluted spray material to be applied to 1000 sq it if turf is matted or certain hard to kill weeds are present, a second application may be required five days later. Area should be dead and ready for seed five days after treatment. Wash sprayer out well after using

Do not remove stubble. Because seed must come in contact with the soil, better results will be obtained if heavily hiatted and bare spots are loosened with an iron rake

Apply a complete fertilizer, such as Gro-Well Lawn Builder according to recommendations for a new lawn

Water and thoroughly soak area prior to seeding

Use a good grade of lawn seed, such as Gro-Wall Premium or Merion Mix and apply.

A very light top dressing of sterilized soil or finely ground - at will improve germination of seed.

Water again lightly in order to wash seed down into soil. Keep area moist with frequent light watering; until at least the first mowing or until new grass fills in. If not and dry, this could require two light waterings daily.

For killing vegetation around trees, shrubs, driveways, patios and other areas use the same dilution rate, thoroughly wetting for age of vegetation to be killed, being most careful not to get any spray material on foliage, stems,

or bank of desirable plants. A repeat treatment may be necessary in five days.

NOTE: Material may crystalize at cold temperatures. This does not harm product. Return to room temperature and shake well

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing or inhalation of spraymist. Wash thoroughly with soap and water after using and before eating or smoking. Keep away from children and benefic as animals. Keep them off treated areas. Do not store near food and feed products. to not use on vegetable gardens. Do not walk from treated areas onto areas which are not to be treated. Do not apply or allow to drift in areas occupied by unprotected persons or animals

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

Licensed for seed bed preparation and scd renovation under Ansul Chemical Co. Palent No. 3,056,668

NET CONTENTS-1 QUART (32 ozs.)

USDA Reg. No. 5535-49

SOLD BY-

L MIKES MALL WIND KINDS

GHOWAL. 8

Flower & Shoot Bed WHIN

6769



CHABBRAGE

grunger.

ACCEPTED

JAN 14 1971

UNDER THE PEDERAL INSECTICIDE FUNGICIDE AND RODENTIAIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO.2535 34. SUBJECT TO ATTACHED COMMENTS.



CHICKNEED

FOR USE IN PREVENTING THE GROWTH OF CERTAIN WEEDS AND GRASSIG IN BEDS OF SHRUBGERY AND ESTABLISHED FLOWERS.

CAPTER HAR IN OF BOOM OF CARD A SECURENCE OF CARD SECURENCE.