

**ACCEPTED**  
 Jun 28 1972  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT,  
 FOR ECONOMIC POISON REGISTERED  
 ED UNDER NO. .... SUBJECT  
 TO ATTACHED COMMENTS.

5535-48

**GRO-WELL** <sup>®</sup>**DYRENE LAWN FUNGICIDE**, works on the leaf surface and when evenly and properly sprayed on lawns will control such diseases as **BROWN PATCH, COPPER SPOT, DOLLAR SPOT, HELMINTHOSPORIUM (Leaf Spot), MELTING OUT, RUST, RED THREAD** and **SNOW MOLD**. For a preventative program begin applications before diseases appear and repeat as needed, usually at 10 to 14 day intervals. If weather is cool and moist then treatments every 7 days may be advisable. This schedule should be followed through the Spring and Summer months.

**FOR CONTROL OF SNOW MOLD (Typhula):** Apply **GRO-WELL** <sup>®</sup>**DYRENE LAWN FUNGICIDE** at 4 ounces per 1,000 sq. ft. after the first killing frost in the Fall. Repeat this application in mid-Winter when no snow is on the ground, following with a third application after the final melting of snow in early Spring.


**FOR THE CONTROL OF THE DISEASES GIVEN ABOVE OTHER THAN SNOW MOLD:** Apply **GRO-WELL** <sup>®</sup>**DYRENE LAWN FUNGICIDE**, at 4 to 8 ounces in 5 gallons of water to 1,000 sq. ft. (20' x 50') repeating treatments at 1 to 2 week intervals being careful to get complete and thorough coverage of the area sprayed. A sprayer which attaches to the hose, such as the **GRO-WELL LAWN SPRAYER**, may be used. Be sure to keep material thoroughly mixed while spraying.

**CAUTION:** Harmful if swallowed or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mix. Wash thoroughly with warm water and soap after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed, foodstuffs, water supplies and fish ponds. Keep out of reach of children. Do not allow spray to drift into areas occupied by unprotected persons or domestic animals. Keep pets and children off treated area until completely dry.

**NOTE:** Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

<sup>®</sup>**DYRENE** is a registered trademark of Farbenfabriken Bayer A. G., Chemagro Corp. licensee.

Sold by: **NET WEIGHT: 14 ounces** USDA Reg. No. 5535-58  
**J. & L. ADIKES, INC., JAMAICA, NEW YORK**



4.49

**PROTECTS YOUR LAWN AGAINST  
 BROWN PATCH, SNOW MOLD,  
 COPPER SPOT, LEAF SPOT, RUST,  
 MELTING OUT & CERTAIN OTHER DISEASES**

<b>ACTIVE INGREDIENTS:</b> 2,4-Dichloro-6-(0-chloroanilino)	By Weight
S-triazine*	50.00%
<b>INERT INGREDIENTS:</b>	50.00%
	<b>Total..... 100.00%</b>

\*U.S. Pat. Nos. 2,720,480 & 3,074,946 — Canadian Pat. No. 546,027  
 This material used under license from Chemagro Corporation.

**COVERS UP TO 3500 SQ. FT.**

**CAUTION:  
 KEEP OUT OF REACH OF CHILDREN  
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

**ACCEPTED**  
 Apr 28 1972  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND MITICIDE ACT  
 THIS LABELING IS SUBJECT  
 TO FEDERAL AND STATE  
 REGULATIONS.

5535-48

**GRO-WELL® DYRENE LAWN FUNGICIDE**, works on the leaf surface and when evenly and properly sprayed on lawns will control such diseases as BROWN PATCH, COPPER SPOT, DOLLAR SPOT, HELMINTHOSPORIUM (Leaf Spot), MELTING OUT, RUST, RED THREAD and SNOW MOLD. For a preventative program begin applications before diseases appear and repeat as needed, usually at 10 to 14 day intervals. If weather is cool and moist then treatments every 7 days may be advisable. This schedule should be followed through the Spring and Summer months.

**FOR CONTROL OF SNOW MOLD** (Typhula): Apply GRO-WELL® DYRENE LAWN FUNGICIDE at 4 ounces per 1,000 sq. ft. after the first killing frost in the Fall. Repeat this application in mid-Winter when no snow is on the ground, following with a third application after the final melting of snow in early Spring.


**FOR THE CONTROL OF THE DISEASES GIVEN ABOVE OTHER THAN SNOW MOLD:** Apply GRO-WELL® DYRENE LAWN FUNGICIDE, at 4 to 8 ounces in 5 gallons of water to 1,000 sq. ft. (20' x 50') repeating treatments at 1 to 2 week intervals being careful to get complete and thorough coverage of the area sprayed. A sprayer which attaches to the hose, such as the GRO-WELL LAWN SPRAYER, may be used. Be sure to keep material thoroughly mixed while spraying.

**CAUTION:** Harmful if swallowed or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mix. Wash thoroughly with warm water and soap after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed, foodstuffs, water supplies and fish ponds. Keep out of reach of children. Do not allow spray to drift into areas occupied by unprotected persons or domestic animals. Keep pets and children off treated area until completely dry.

**NOTE:** Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

©DYRENE is a registered trademark of Farbenfabriken Bayer A. G., Chemagro Corp. licensee.

Sold by: **NET WEIGHT: 14 ounces** USDA Reg. No. 5535-58  
**J. & L. ADIKES, INC., JAMAICA, NEW YORK**



**4.49**

**PROTECTS YOUR LAWN AGAINST  
 BROWN PATCH, SNOW MOLD,  
 COPPER SPOT, LEAF SPOT, RUST,  
 MELTING OUT & CERTAIN OTHER DISEASES**

<b>ACTIVE INGREDIENTS:</b> 2,4-Dichloro-6-(0-chloroanilino)	By Weight
S-triazine*	50.00%
<b>INERT INGREDIENTS:</b>	50.00%
	<b>Total.....100.00%</b>

\*U.S. Pat. Nos. 2,720,480 & 3,074,946 — Canadian Pat. No. 546,027  
 This material used under license from Chemagro Corporation.

**COVERS UP TO 3500 SQ. FT.**

**CAUTION:  
 KEEP OUT OF REACH OF CHILDREN  
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS**