

GRASS-WEED KILLER is a water soluble powder which is used to control certain diseases on LAWNS, OPEN SPACES, FRUIT and VEGETABLES. Unless otherwise instructed, apply sprays every 7-10 days, beginning when diseases are first apparent and continuing as long as disease conditions are present. Cover thoroughly, but not in excess, being sure to spray underside of leaves. A sprayer which attaches to a hose, such as the GRASS-WEED KILLER SPRAYER, may be used to apply, except on lawns, where the GRASS-WEED KILLER SPRAYER should be used.

DO NOT USE ON FRUIT WITH 1 GAL. OF LIQUID EXCEPT LAWNS (SEE NOTE 1 BELOW)

FOR USE ON	SEE NOTE	TO CONTROL	DOSAGE (Application Rate)
LAWNS: Bluegrass, Fescue, Ryegrass, Bentgrass	1-4	Leaf Spot (Helminthosporium)	2 ozs. (12 tbsps.)
Bluegrass (Merion)	1-2	Rust	
Bluegrass, Fescue, Ryegrass, Bentgrass, Bermuda grass	1-3	Pythium Blight	
TREES-SHRUBS: Sycamore	5	Anthracnose	4 tps.
Oak	6		
Dogwood	7	Blight (Flower & Leaf)	1 tbsp.
Azalea	8	Calls (Azalea & Pinkster)	4 tbsps.
Azalea, Camellia	9	Fatal Blight	2 tbsps.
FRUITERS: Apples	10	Leaf Spot	
Gladiolus	10-11	Potrytis Blight, Curvularia, Sterchylium	
Carnation, Chrysanthemum, Hollyhock, Spanghann	12	Anthracnose, Leaf Spot, Rust	
FRUITERS: Apples	13	Blotch (Early & Late), Procks Spot, Cedar Rust, Froeye, Rot (Black & Bitter), Leaf	
Apples (Do not apply within 20 days of harvest)	14	Fire Blight	
Pears (Do not apply within 7 days of harvest)			
Peaches			
VEGETABLES: Broccoli, Cabbage, Cauliflower (Do not apply within 7 days of harvest)		Alternaria, Downy Mildew	
Beans (Do not apply within 7 days of harvest)	16	Anthracnose, Downy Mildew, Rust	
Cantaloupe, Melons, Cucumbers, Squash, Watermelon (Do not apply within 7 days of harvest)	17	Alternaria (Main species), Blight, Anthracnose, Downy Mildew, Early Stem Blight	
Tomatoes (Do not apply within 7 days of harvest)		Anthracnose, Blight (Early & Late), Leaf Spot, Verticillium	

NOTES: (1) Dilute with 5 gals. of water and apply to 1,000 sq. ft. (20'x50'). Do not graze treated areas and do not feed clippings to livestock. (2) Apply every 7-10 days when disease is evident, usually during July to October. (3) Apply every 5-10 days from July to September. (4) Apply every 7-14 days during July and August. (5) Make 2 applications: the first as buds start to swell; second when buds are open. Third if necessary later. As a special precaution, remove dead leaves in the fall. (6) Do not apply to plants which are in bloom. (7) 10-14 days later and re-treat if necessary. (8) Do not apply to plants which are in bloom. (9) Apply at bud-break, re-treat if necessary. (10) Do not apply to plants which are in bloom. (11) Do not apply to plants which are in bloom. (12) Do not apply to plants which are in bloom. (13) Do not apply to plants which are in bloom. (14) Do not apply to plants which are in bloom. (15) Do not apply to plants which are in bloom. (16) Do not apply to plants which are in bloom. (17) Do not apply to plants which are in bloom.

DO NOT USE ON FRUIT WITH 1 GAL. OF LIQUID EXCEPT LAWNS (SEE NOTE 1 BELOW)

GROWELL

LAWN FUNGICIDE CONTAINS

DYRENE[®]

Prevents and Controls
Lawn Diseases &
Brown Patch, Dollar
& Leaf Spot, Melting Out,
Rust, Snow Mold

ACCEPTED

MAR 31 1983

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Keep out of reach of children. Do not breathe dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco, or using the toilet. Wash hands before eating, drinking, using tobacco, or using the toilet. Do not get in eyes, on clothing, or on food. Do not use near water supplies and fish ponds. Do not use in areas where children and domestic animals play. Do not use in areas where livestock graze.

ENVIRONMENTAL HAZARDS

Do not use in areas where waterfowl, wild birds, or other wildlife are present. Do not use in areas where bees are present. Do not use in areas where fish are present. Do not use in areas where livestock graze.

DIRECTIONS FOR USE

Apply to lawn as directed. Do not use on other plants. Do not use on lawns that are dormant. Do not use on lawns that are heavily shaded. Do not use on lawns that are heavily mowed. Do not use on lawns that are heavily fertilized.

STORAGE

Store in original container. Do not use if container is damaged. Do not use if contents are solid or if odor is present.

RENTAL/REUSE

Do not use if container is damaged. Do not use if contents are solid or if odor is present.

CONTAINER DISPOSAL

Do not reuse. Do not burn. Do not dump in water.

ASSURANCE STATEMENT

This product has been registered with the Environmental Protection Agency. It meets the requirements of the Federal Insecticide, Fungicide, and Rodenticide Act.

Manufactured by Growell Chemical Company, Inc., 10000 N. 10th Ave., Minneapolis, MN 55412.

© 1983 Growell Chemical Company, Inc. All rights reserved.

NET WEIGHT 1.5 LBS (680g) 100% Active Ingredient

DOSE: 1/2 lb per 1000 sq ft

Apply to lawn as directed. Do not use on other plants.

Do not use on lawns that are dormant. Do not use on lawns that are heavily shaded.

Do not use on lawns that are heavily mowed. Do not use on lawns that are heavily fertilized.