


FRONT
PANEL




1.49

Supplies - 1967

INSECTICIDE
FUNGICIDE
MITICIDE
SINCE 1855

ACCEPTED
DEC 14 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT GIDE ACT
FOR ECONOMIC POISON REG. STR.
ED UNDER NO. 2229-66 SUBJECT
TO ATTACHED COMMENTS.



CONTAINS:
MALATHION*
SEVIN** & PHALTAN***

Faint, illegible text at the bottom of the page.

[282]

REAR
PANEL

Camerman's Note
Poor Copy

GRO-WELL ROSE DUST is an Insecticide-Fungicide-Miticide combination that is light in color and will therefore not leave unsightly dark residues on Rose petals or the blooms of other flowering plants.

To be most effective, GRO-WELL ROSE DUST should be applied before insects become numerous and before diseases are apparent. Protect plants through the growing season by dusting at 7 to 14 day intervals and after heavy rains. Cover foliage thoroughly, but do not use in excess. Pay particular attention to cover underside of leaves.

DIRECTIONS:


Lift hinged cap and squeeze container to dust. Squeeze Duster may be refilled from the 1 lb. or 4 lb. can of GRO-WELL FLORAL & ROSE DUST.

INSECTS

Effective for the control of the following insects: Aphids, Cicada (17 year Locusts), Clover Mites, Red Spider Mites, Japanese Beetles, Rose Chafer, Rose Leathoppers, Leaf Rollers, Rose Slugs, Flea Beetles, Turps, Tarnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on Roses, Gladiolus, Dahlias, Sweet Peas, Asters, Chrysanthemums, Delphinium, Calendula, Begonia and Snapdragon.

DISEASES

Black Spot and Powdery Mildew of Roses, Septoria, Leaf Spot of Chrysanthemums, Didymella, Leaf Spot of Iris, Rust and Alternaria Leaf Spot of Carnations, Alternaria Leaf Spot, Powdery Mildew and Anthracnose on Zinnias, Marigolds, Asters, Phlox and Snapdragons.



CAUTION: Keep out of reach of children and pets. Harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water: for eyes, get medical attention. Wash hands thoroughly before eating or smoking. Do not contaminate feed and foodstuffs. Do not apply or allow to drift into areas occupied by unprotected persons or beneficial animals. If spilled, flush with plenty of water.

Do not contaminate fish ponds. Do not apply where runoff is likely to occur. Do not use on certain Ferns including Boston, Maidenhair and Pteris, some species of Crassula and Boston Ivy.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS:

*Malathion 4.00%
**Carbaryl (1-Naphthyl N-Methylcarbamate) 3.00%
***Folpet (N-Trichloromethylthio) phthalimide 5.00%

INERT INGREDIENTS 88.00%

*O,O-dimethyl-dithiophosphate of diethyl mercaptosuccinate.
**Registered Trade Mark of Union Carbide Corp. for Carbaryl Insecticide.
***Phaltan is registered Trade Mark for Folpet Fungicide. U.S. Pat. Nos. 2,553,770 - 2,553,771.

NET WEIGHT 10 OUNCES

SOLD BY

U.S.D.A. Reg. No. 5535-

GENK S. R.L.

[282]

REAR
PAGE

GRO-WELL ROSE DUST is an Insecticide-Fungicide Miticide combination that is light in color and will therefore not leave unightly dark residues on Rose petals or the blooms of other flowering plants.

To be most effective, GRO-WELL ROSE DUST should be applied before insects become numerous and before diseases are apparent. Protect plants through the growing season by dusting at 7 to 14 day intervals and after heavy rains. Cover foliage thoroughly, but do not use in excess. Pay particular attention to cover underside of leaves.

DIRECTIONS:

Lift hinged cap and squeeze container to dust. Squeeze Duster may be refilled from the 1 lb. or 4 lb. can of GRO-WELL FLORAL & ROSE DUST.

INSECTS

Effective for the control of the following insects: Aphids, Cicada (17 year Locusts), Clover Mites, Red Spider Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Flea beetles, Tangle-farnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on Roses, Gladiolus, Dahlias, Sweet Pea, Asters, Chrysanthemums, Delphinium, Calendula, Begonia and Snapdragon

DISEASES

Black Spot and Powdery Mildew of Roses, Septoria, Leaf Spot of Chrysanthemums, Didymellina, Leaf Spot of Iris, Rust and Alternaria Leaf Spot of Carnations, Alternaria Leaf Spot, Powdery Mildew and Anthracnose on Zinnia, Impatiens, Asters, Philoxera and Scab on geranium

ACCEPTED
Dec. 14, 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE ACT
REGISTERED UNDER NO. 5535-16
SUBJECT TO ATTACHED COMMENTS.

CAUTION: Keep out of reach of children and pets. Avoid contact with eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. In case of contact, flush with water. For eyes, get medical attention. Wash hands thoroughly after use. Do not smoke. Do not contaminate food and foodstuffs. Do not apply directly to

CAUTION: Keep out of reach of children and pets. Harmful if swallowed. May cause irritation of nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water: for eyes, get medical attention. Wash hands thoroughly before eating or smoking. Do not contaminate feed and foodstuffs. Do not apply or allow to drift into areas occupied by unprotected persons or beneficial animals. If spilled, flush with plenty of water.

Do not contaminate fish ponds. Do not apply where runoff is likely to occur. Do not use on certain Ferns including Boston, Maidenhair and some species of Crassula and Boston Ivy.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS:

*Malathion	4.00%
**Carbaryl (1-Naphthyl N-Methylcarbamate)	3.00%
***Folpet (N-Trichloromethylthio) phthalimide	5.00%

INERT INGREDIENTS 88.00%

- *O,O-dimethyl-dithiophosphate of diethyl mercaptosuccinate.
- **Registered Trade Mark of Union Carbide Corp. for Carbaryl Insecticide.
- ***Phaltan is registered Trade Mark for Folpet Fungicide. U.S. Pat. Nos. 2,553,770 - 2,553,771.

NET WEIGHT 10 OUNCES

SOLD BY

U.S.D.A. Reg. No. 5535