


FRONT
PANEL



ACCEPTED
JAN 11 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 22,322-67 SUBJECT
TO ATTACHED COMMENTS.

[292]

CRIVELLI MULTI-PURPOSE DUST is effective for control of the following insects and diseases on FRUIT TREES, SHRUBS, BUCKS and VEGETABLES:

INSECTS:
 Aphids, Beet Beetles, Borers, Caterpillars, Flea Beetles, Fruit Flies, Fruit Weevils, Greenhouse Aphids, Leaf Beetles, Leaf Miners, Leafhoppers, Mosquitoes, Nymphs, Sawflies, Scale Insects, Spiders, Thrips, Whiteflies, Yellowjackets.

DISEASES:
 Anthracnose, Blight, Botrytis, Brown Rot, Fire Blight, Fruit Rot, Mildew, Powdery Mildew, Rust, Scab, Verticillium Wilt, Xanthomonas.

VEGETABLES: Beans, Broccoli, Cabbage, Cauliflower, Cucumbers, Eggplant, Kale, Lettuce, Melons, Peas, Potatoes, Spinach, Squash, Tomatoes, Zucchini.

FRUIT TREES: Apples, Citrus, Grapes, Peaches, Plums, Pears, Prunes, Raspberries, Strawberries, Vines.

SHRUBS: Camellias, Hibiscus, Magnolias, Roses, Spirea, Yucca.

BUCKS: Deer, Rabbits, Squirrels.

ACCEPTED

JAN 11 1968

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PESTICIDE REGISTERED UNDER NO. 5539-67 SUBJECT TO ATTACHED COMMENTS.

CRIVELLI MULTI-PURPOSE DUST is effective for control of the following insects and diseases on FRUIT TREES, SHRUBS, BUCKS and VEGETABLES:

INSECTS:
 Aphids, Beet Beetles, Borers, Caterpillars, Flea Beetles, Fruit Flies, Fruit Weevils, Greenhouse Aphids, Leaf Beetles, Leaf Miners, Leafhoppers, Mosquitoes, Nymphs, Sawflies, Scale Insects, Spiders, Thrips, Whiteflies, Yellowjackets.

DISEASES:
 Anthracnose, Blight, Botrytis, Brown Rot, Fire Blight, Fruit Rot, Mildew, Powdery Mildew, Rust, Scab, Verticillium Wilt, Xanthomonas.

VEGETABLES: Beans, Broccoli, Cabbage, Cauliflower, Cucumbers, Eggplant, Kale, Lettuce, Melons, Peas, Potatoes, Spinach, Squash, Tomatoes, Zucchini.

FRUIT TREES: Apples, Citrus, Grapes, Peaches, Plums, Pears, Prunes, Raspberries, Strawberries, Vines.

SHRUBS: Camellias, Hibiscus, Magnolias, Roses, Spirea, Yucca.

BUCKS: Deer, Rabbits, Squirrels.

0

[293]

ONE-STEP MULTI-PURPOSE DUST is effective for control of the following INSECTS and DISEASES on ROSES, FRUIT TREES, SHRUBS, HERBS and VEGETABLES.

INSTRUCTIONS:

Put large cap and squeeze container to dust. Squeeze Duster may be rented from the 1 lb. or 4 lb. can of ONE-STEP MULTI-PURPOSE DUST. Apply to Roses, Fruit Trees, Shrubs and Vegetables before insects become numerous and before diseases become apparent. Protect plants through the growing season by dusting at 7 to 14 day intervals and after heavy rains. Cover foliage thoroughly, but do not use in excess. Pay particular attention to cover underside of leaves. On Fruit Trees, apply the material directly to the tree when the bark is still. For timing of applications, follow Fruit Tree spray and dust schedule.

ROSES: Aphids, Spider Mites, Japanese Beetles, Rose Chafers, Rose Leafhoppers, Leaf Beetles, Rose Stems, Flea Beetles, Thrips, Tarnished Plant Hoppers, Four-lined Leaf Beetles, Whitefly, Asiatic Garden Beetle and Black Blister Beetle, on Roses, Gladiolus, Dahlias, Sweet Peas, Aster and Chrysanthemums, Black Spot, Powdery Mildew, Rust, Leaf Spot and Botrytis, Flower blight of Roses, Tuberosa Legumes, Clematis, Chrysanthemum and Peonies.

FRUIT TREES: Aphids, Colling Moth, Nates (European Red, Clover, Two Spotted, White-necked), Pear Psylla, Japanese Beetles, Black Cherry Aphid, Tent Caterpillar, Apple Sawb, Sooty Blotch, Brooks Spot, Winter Rot, Fly Speck, Brown Rot, Peach Scab, Leaf Spot on Apples, Pears, Potatoes and Cherries.

SHRUBS: Aphids, Spider Mites, Lacebugs, Scale Insects, Scurfy, Leafminers, Beetles, Caterpillars, Bagworms and Boxwood Leaf Miner on Azaleas, Rhododendron, Magnolias, Lilacs, Hydrangeas, Euonymus, Fuchsia Japonica, Hibiscus and Boxwood. Note: For scale insects and bagworms on Surtida, make two dusting applications. Do not spray for control of these insects as 2 or 3 times the normal dosage of this product is required per gallon of water. This would result in excessive blighting of spray nozzles. For many diseases on Hibiscus and Euonymus.

HERBS: Aphids, Spider Mites, Flea Beetles, Leafminers, Caterpillars and Aphids on Parsnips, Potatoes and Peas.

VEGETABLES: Spider Mites, Flea Beetles, Leafminers, Caterpillars, European Pine Sawb, Tent Caterpillars and Aphids on Cabbages, Peaches, Nuts, Beans, Cucumbers, Tomatoes and Potatoes.

FRUIT TREES: Aphids, Spider Mites, Flea Beetles, Colorado Potato Beetle, early and late blight, Anthracnose and Septoria on Potatoes.

VEGETABLES: Aphids, Flea Beetles, Colorado Potato Beetle and Leafminers on Potatoes.

FRUIT TREES: Aphids, Colorado Potato Beetle, Flea Beetles, Leafminers, Scurfy and Leaf Blight.

VEGETABLES: Aphids, Colorado Potato Beetle, Flea Beetles, Leafminers, Scurfy and Leaf Blight.

FRUIT TREES: Aphids, Colorado Potato Beetle, Flea Beetles, Leafminers, Scurfy and Leaf Blight.

VEGETABLES: Aphids, Colorado Potato Beetle, Flea Beetles, Leafminers, Scurfy and Leaf Blight.

FRUIT TREES: Aphids, Colorado Potato Beetle, Flea Beetles, Leafminers, Scurfy and Leaf Blight.

VEGETABLES: Aphids, Colorado Potato Beetle, Flea Beetles, Leafminers, Scurfy and Leaf Blight.

ACCEPTED

JAN 11 1968
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 53267 SUBJECT TO ATTACHED COMMENTS.

animals. If spilled, flush area with plenty of water.

Do not use Sulphur on Apples within 10 days to 2 weeks of an oil spray.

Do not make applications when temperature is high as Sulphur may burn foliage.

Do not apply to Red Delicious Apples as the combination of Captan and Sulphur

may cause foliage injury. Injury may occur on Rose Pears in the Northeast

under certain conditions. Injury may occur to certain varieties of Sweet

Cherries, especially in the Northwest. Do not apply to Apples, cherries,

Eggplants and Peas within 3 days of harvest. Do not apply to Peas within

1 day of harvest. Do not apply to Tomatoes within 5 days of harvest. Do

not apply to Peaches within 7 days of harvest. After the time limits set

above have passed, an effective fungicide should be used alone in a manner

which would not leave residues in excess of established tolerances. Do not

use on crops to be fed to dairy animals or to animals being finished for

slaughter.

Do not contaminate fish ponds. Do not apply when snow is likely to occur.

NOTE: Follow directions carefully, as use, storage and handling of this

material is beyond the control of the seller. The buyer assumes all liability.

Directions given hereon are not strictly adhered to.

ACTIVE INGREDIENTS:

Captan	40%
Captan (1,1-dichloro-3,3-dimethyl-2-butylthiourea)	50%
Captan (1,1-dichloro-3,3-dimethyl-2-butylthiourea)	40%
Sulphur	25.00%

Half Inert Ingredients 62.00%

100% dimethyl diethyl malonate or dimethyl mercaptosuccinate.
*Sulphur is Reg. U.S. Pat. & Trad. Office, U.S. Chem. & Eng. Div., Wash., D.C.

U.S. Pat. 2,811,100

U.S. Pat. 2,811,100

U.S. Pat. 2,811,100