


5535-75

ACCEPTED
 JUN 30 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR FURNISHING TO THE PUBLIC
 TO ATTACHED



2M-11-70

CONTROL
 GYPSY MOTH, INCHWORMS, JAPANESE BEETLES and
 CERTAIN OTHER INSECTS ON
 ROSES, FLOWERS, SHRUBS, TREES, FRUITS AND VEGETABLES

ACTIVE INGREDIENTS:	
*Methoxychlor, Technical	25.40%
Xylene	71.60%
INERT INGREDIENTS:	3.00%
TOTAL	100.00%

*Equivalent to 22.352% 2,2-bis(p-methoxyphenyl)-1,1,1-Trichloroethane and 3.048% other isomers and reaction products.

**CAUTION: KEEP OUT OF REACH OF CHILDREN
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

MAKES UP TO 24 GALLONS OF SPRAY


Read entire label. Use strictly in accordance with label cautions, warnings and directions. Do not store diluted spray. Make a new dilution for each use. Spray thoroughly, wetting all leaf surfaces, including the underside, before insects become numerous. Spray at weekly intervals as needed.

DOSAGE LISTED IS TO BE MIXED WITH 1 GALLON OF WATER

ROSE-FLOWERS EVERGREENS-SHRUBS	INSECTS	DOSAGE	SHADE TREES- ORNAMENTALS	INSECTS	DOSAGE
Andromeda, Arbutus, Asters, Azaleas, Begonias, Berwood, Camellias, Carnations, Chrysanthemums, Coleus, Dahlias, Delphiniums, Eranthis, Juniper, Marigolds, Nasturtiums, Rhododendron, Roses, Snap, Dragon, Yucca, Zinnias.	Eyes, Blister and Jap Spotted; Leaf & Flea Hoppers, Licebugs, Caterpillars, Rose Chafar & Rose Slugs.	4 to 6 tps.	Birch, Dogwood, Elm, Holly, Magnolia, Maple, Oak, Pine and Sycamore.	Centipedes, Gypsy and Tussock Moth, Insects, Juniper Webworm and Pine Sawflies. Scale Insects and Elm Bark Beetle. Do not apply.	4 to 6 tps. 8 to 12 tps.
Do not use on Petunias, Boston Maiden Hair, Pteris Fern and Crassula.			Do not spray Chinese Elms, Jap Maples, Red Maples and Red-bud as foliage may be injured.		
FRUITS	INSECTS	DOSAGE	VEGETABLES	INSECTS	DOSAGE
Apples, Cherries, Pears and Plums—Do not apply within 7 days of harvest. Apricots & Peaches—Do not apply within 21 days of harvest. Blackberries, Blueberries, Currants, Gooseberries, Grapes, Raspberries and Strawberries—Do not apply within 14 days of harvest.	Apple Maggot, Cherry Fruit Worm, Codling Moth, Leaf Miner, Plum Curculio, Tent Caterpillar and Saw Beetsle.	4 to 6 tps.	Beans, Cabbage, Cauliflower, Eggplant, Peas, Peppers, Potatoes and Tomatoes—Do not apply within 7 days of harvest.	Beetle, Bean Leaf, Colorado Potato Beetle, Flea and Jap. WORMS: Cabbage, Fall Army, Tomato Fruit, Green Horn and Web; and Lacebugs.	4 to 6 tps.
	Caterpillar, Cherry Fruit Fly, Grape Berry Moth, Grape Leaf Stipeworm, Oriental Fruit Moth, Pear Chafar, Leaf Miner, Spittlebug. See also Scale Crawler and Pear Slug.	4 to 6 tps.			

NOTE: DO NOT spray when temperature is above 85°F. Do not use on crops within 14 days of application of any sulphur product. Wash residue off all fruits and vegetables before using.
WARNING: HARMFUL IF SWALLOWED. Keep out of reach of children. Do not inhale spray mist. Avoid skin contact. If contacted wash immediately with soap and water. Do not use on pets, humans or animals. Do not apply or allow to drift into areas occupied by unprotected persons or domestic animals. Wash well after using and before smoking or eating. Do not use on forage crops fed to dairy animals. Do not contaminate feed or foodstuffs. Do not store or use near fire or open flame. Dispose of this container when empty. Do not use for any other purpose.
 Keep out of lakes, streams or ponds.
 This product is toxic to bees and should not be applied when bees are active in the area.
NOTICE: Follow directions carefully. As use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

SOLD BY **NET CONTENTS 16 OUNCES** U.S.D.A. Reg. No. 5535-75
J. & L. ADIKES, INC., JAMAICA, NEW YORK 11423



- PRODUCTS**
- MULTI-PURPOSE DUST
 - LAWN FERTILIZER
 - LAWN WEED KILLER
 - GRASS KILLER
 - SPRING FERTILIZER
 - ROSE ARBORAL SPRAY
 - CONCRETE SPRAY
 - FRUIT TREE SPRAY
 - ROSE DUST
 - TOMATO & VEGETABLE DUST
 - INSECTICIDE
 - TREE WOUND PAINT
 - HONEY DEW
 - GRASS KILLER
 - GRO-WELL LAWN SEED
 - LAWN FOOD
 - ACID PLANT FOOD
 - GARDEN PLANT FOOD
 - ROSE FOOD

FOR
ED UNDER
TO ATTACHED

2.98

GRO-WELL METHOXYCHLOR will control the insects listed below which attack Roses, Shrubs, Trees, Evergreens, Fruit Trees and Vegetables. A sprayer which attaches to a hose, such as the GRO-WELL SHRUB & TREE SPRAYER, may be used. Read entire label. Use strictly in accordance with label cautions, warnings and directions. Do not store diluted spray. Make a new dilution for each use. Spray thoroughly, wetting all leaf surfaces, including the underside, before insects become numerous. Spray at weekly intervals as needed.

DOSAGE LISTED IS TO BE MIXED WITH 1 GALLON OF WATER

ROSE-FLOWERS EVERGREENS-SHRUBS	INSECTS	DOSAGE	SHADE TREES— ORNAMENTALS	INSECTS	DOSAGE
Andromedas, Arborvitae, Asters, Azaleas, Begonias, Boxwood, Camellias, Carnations, Chrysanthemums, Coleus, Dahlias, Delphiniums, Euonymous, Geraniums, Gladioli, Juniper, Marigolds, Nasturtiums, Rhododendron, Roses, Snap, Dragons, Taxus, Zinnias.	Flea, Blister and Jap Beetles; Leaf & Flea Hoppers, Lacebugs, Cankerworms, Rose Chafer & Rose Slugs.	4 to 6 tsps.	Birch, Dogwood, Elm, Holly, Magnolia, Maple, Oak, Pine and Sycamore.	Cankerworms, Gypsy and Tussock Moth, Inchworm, Juniper Webworm and Pine Sawflies. Scale Insects and Elm Bark Beetle. Dormant Application only.	4 to 6 tsps. 8 to 12 tsps.
Do not use on Petunias, Boston Maiden Hair, Pteris Fern and Crassula.			Do not spray Chinese Elms, Jap Maples, Red Maples and Red-bud as foliage may be injured.		
FRUITS	INSECTS	DOSAGE	VEGETABLES	INSECTS	DOSAGE
Apples, Cherries, Pears and Plums—Do not apply within 7 days of harvest.	Apple Maggot, Cherry Fruit Worm, Codling Moth, Leaf Hopper, Plum Curculio, Tent Caterpillar and Jap Beetle.	4 to 6 tsps.	Beans, Cabbage, Cauliflower, Eggplant, Peas, Peppers, Potatoes and Tomatoes—Do not apply within 7 days of harvest.	Beetles, Bean Leaf, Blister, Colorado Potato, Mexican Bean, Flea and Jap; WORMS: Cabbage, Fall Army, Tomato Fruit, Green Horn and Web; and Lacebugs.	4 to 6 tsps.
Apricots & Peaches—Do not apply within 21 days of harvest.					
Blackberries, Blueberries, Currants, Gooseberries, Grapes, Raspberries and Strawberries—Do not apply within 14 days of harvest.	Cankerworm, Cherry Fruit Fly, Grape Berry Moth, Grape Leaf Skeletonizer, Oriental Fruit Moth, Rose Chafer, Leaf Hopper, Spittlebug, San Jose Scale Crawler and Pear Slug.	4 to 6 tsps.			


NOTE: DO NOT spray when temperature is above 85°F. Do not use on crops within 14 days of application of any sulphur product. Wash residue off all fruits and vegetables before using.

WARNING: HARMFUL IF SWALLOWED. Keep out of reach of children. Do not inhale spray mist. Avoid skin contact. If contacted wash immediately with soap and water. Do not use on pets, humans or animals. Do not apply or allow to drift into areas occupied by unprotected persons or domestic animals. Wash well after using and before smoking or eating. Do not use on forage crops fed to dairy animals. Do not contaminate feed or foodstuffs. Do not store or use near fire or open flame. Dispose of this container when empty. Do not use for any other purpose. Keep out of lakes, streams or ponds.

This product is toxic to bees and should not be applied when bees are active in the area.
NOTICE: Follow directions carefully. As use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

SOLD BY **NET CONTENTS 16 OUNCES** U.S.D.A. Reg. No. 5535-75

J. & L. ADIKES, INC., JAMAICA, NEW YORK 11423



PRODUCTS

- MULTI-PURPOSE DUST
- LAWN FUNGICIDE
- LAWN WEED KILLER
- CRACKGRASS KILLER
- INDOOR-OUTDOOR AEROSOL SPRAY
- ROSE AEROSOL SPRAY
- DORMANT SPRAY
- FRUIT TREE SPRAY
- ROSE DUST
- TOMATO & VEGETABLE DUST
- MALATHION
- TREE WOUND PAINT
- HORNET BOMB
- CHICKWEED KILLER
- GRO-WELL LAWN SEED
- LAWN FOOD
- ACID PLANT FOOD
- GARDEN PLANT FOOD
- ROSE FOOD