

## GRO-WELL 5% SEVIN\* DUST

**LAWNS AND FOUNDATION PLANTINGS:** For control of Ants, Chinch Bugs, Lawn Moths, (Sod Webworms), Cutworms, Earwigs, Ticks, Centipedes, Grass Hoppers and Armyworms, apply 1 pound evenly to an area of 250 sq. ft. (10' x 25').

**ORNAMENTALS:** Dust when insects appear and repeat as necessary to control Japanese and Flea Beetles, Bagworm, Boxwood Leaf Miner, Leaf Rollers, Periodical Cicada, Rose Aphid, Rose Slug, Scale insects and Thrips. (NOTE: Do not use on Boston Ivy. For optimum scale control apply when crawlers are present.)

**VEGETABLES: BEANS**—Mexican Bean Beetle, Leaf Hoppers, Lygus and Stink Bugs, use 1/2 to 1 lb. per 1,000 sq. ft. (50' x 20'); Bean Leaf, Cucumber, Flea and Japanese Beetles use 1 lb. per 1,000 sq. ft. **CUCUMBERS AND SUMMER SQUASH**—Pickleworm, Melonworm, Cucumber and Flea Beetles use 1/2 lb. per 1,000 sq. ft. **TOMATO, EGGPLANT AND PEPPER**—Tomato Fruitworm and Hornworm, Fall Armyworm, Flea, Beetle, Leaf Hoppers, European Corn Borer and Stink Bugs use 1/2 to 1 lb. per 1,000 sq. ft. **POTATO**—Colorado Potato and Flea Beetles and Leaf Hoppers use 1/2 to 1 lb. per 1,000 sq. ft. **CORN**—Corn Earworm, European Corn Borer, Fall Armyworm, Sap and Japanese Beetles use 1 lb. per 1,000 sq. ft. (NOTE: For Corn Earworm on Corn make 3 or more applications to the silks at two to three day intervals, starting when silks first appear. For European Corn Borers make thorough application into the whorls when plant is knee high and repeat at weekly intervals until ears are ready for picking.)

**BULBS:** This product may be used as a bulb dust to control thrips while bulbs are in storage. Put 1 tablespoon of this material in a paper bag and put approximately 50 bulbs in bag and shake bag until all bulbs are coated with dust.

### CAUTION:

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid contact with skin, eyes and clothing. Avoid contact. Wash hands with soap and warm water before eating or smoking. Do not reuse container. Destroy when empty.

May be applied up to and including the day of harvest on crops listed above for human consumption. Do not apply within 7 days of harvest for forage. Do not feed treated bean vines to meat and dairy animals.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

**NOTE:** Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

\*Trade mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

SOLD BY **J. & L. ADIKES, INC., JAMAICA, NEW YORK 11423** NET CONTENTS 1 LB. U.S.D.A. REG. No. 5535 82



**KILLS ANTS,  
CHINCH BUGS,  
CUTWORMS,  
EARWIGS  
and  
CERTAIN OTHER  
INSECTS**



MAY ALSO BE USED AS A BULB DUST

ACTIVE INGREDIENTS:	SINCE 1888	
Carbaryl 1-naphthyl N-methylcarbamate		5.00%
INERT INGREDIENTS		95.00%
	TOTAL	100.00%

COVERS UP TO 2000 SQ. FT.

### CAUTION:

KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

## GRO-WELL 5% SEVIN\* DUST

**LAWNS AND FOUNDATION PLANTINGS:** For control of Ants, Chinch Bugs, Lawn Moths, (Sod Webworms), Cutworms, Earwigs, Ticks, Centipedes, Grass Hoppers and Armyworms, apply 1 pound evenly to an area of 250 sq. ft. (10' x 25').

**ORNAMENTALS:** Dust when insects appear and repeat as necessary to control Japanese and Flea Beetles, Bagworm, Boxwood Leaf Miner, Leaf Rollers, Periodical Cicada, Rose Aphid, Rose Slug, Scale insects and Thrips. (NOTE: Do not use on Boston Ivy. For optimum scale control apply when crawlers are present.)

**VEGETABLES: BEANS**—Mexican Bean Beetle, Leaf Hoppers, Lygus and Stink Bugs, use 1/2 to 1 lb. per 1,000 sq. ft. (50' x 20'); Bean Leaf, Cucumber, Flea and Japanese Beetles use 1 lb. per 1,000 sq. ft. **CUCUMBERS AND SUMMER SQUASH**—Pickleworm, Melonworm, Cucumber and Flea Beetles use 1/2 lb. per 1,000 sq. ft. **TOMATO, EGGPLANT AND PEPPER**—Tomato Fruitworm and Hornworm, Fall Armyworm, Flea Beetle, Leaf Hoppers, European Corn Borer and Stink Bugs use 1/2 to 1 lb. per 1,000 sq. ft. **POTATO**—Colorado Potato and Flea Beetles and Leaf Hoppers use 1/2 to 1 lb. per 1,000 sq. ft. **CORN**—Corn Earworm, European Corn Borer, Fall Armyworm, Sap and Japanese Beetles use 1 lb. per 1,000 sq. ft. (NOTE: For Corn Earworm on Corn make 3 or more applications to the silks at two to three day intervals, starting when silks first appear. For European Corn Borers make thorough application into the whorls when plant is knee high and repeat at weekly intervals until ears are ready for picking.)

**BULBS:** This product may be used as a bulb dust to control thrips while bulbs are in storage. Put 1 tablespoon of this material in a paper bag and put approximately 50 bulbs in bag and shake bag until all bulbs are coated with dust.

### CAUTION:

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid contact with skin, eyes and clothing. Avoid contact. Wash hands with soap and warm water before eating or smoking. Do not reuse container. Destroy when empty.

May be applied up to and including the day of harvest on crops listed above for human consumption. Do not apply within 7 days of harvest for forage. Do not feed treated bean vines to meat and dairy animals.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

**NOTE:** Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

\* Trade mark of Union Carbide Corporation for 1 naphthyl N methylcarbamate

SOLD BY

NET CONTENTS 1 LB.

U.S.D.A. REG. No. 5535-82

# GRO-WELL

98c

## 5%

# SEVIN DUST

KILLS ANTS,  
CHINCH BUGS,  
CUTWORMS,  
EARWIGS

and

CERTAIN OTHER  
INSECTS



MAY ALSO BE USED AS A BULB DUST

### ACTIVE INGREDIENTS:

Carbaryl 1 naphthyl N methylcarbamate

### INERT INGREDIENTS

SINCE 1855

TOTAL

5.00%

95.00%

100.00%

COVERS UP TO 2000 SQ. FT.

CAUTION:

KEEP OUT OF REACH OF CHILDREN.  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS.