





ACCEPTED
July 3, 1971
UNDER THE FEDERAL INSECTICIDE ACT
FOR USE IN THE UNITED STATES
AND POSSESSOR REGIS. NO. 5535-83
EPA REG. NO. 5535-83



ACCEPTED FOR MAILING
 Aug 3, 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND MITTICIDE ACT
 AND ECONOMIC POISON REGISTER
 ED UNDER NO. _____ SUBJECT
 TO ATTACHED LABELS
 55 35 2



CHINCH
BUGS



LAWN MOTHS
(Sod Webworm
Adults)



SOD WEBWORMS
(Lawn Moth
Larvae)

AND
CERTAIN OTHER
ANNOYING
LAWN
INSECTS

KEEP OUT OF REACH OF CHILDREN
 SEE BACK LABEL FOR ADDITIONAL CAUTIONS

INERT INGREDIENTS:

SOLD BY:

NET WEIGHT 7 LBS.

U.S. D. A. Reg. No. [REDACTED]

100 A. ANTHER... [REDACTED]

ACCEPTED
July 3, 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND MITICIDE ACT
 FOR ECONOMIC PESTICIDE REGISTER-
 ED UNDER NO. [REDACTED] SUBJECT
 TO ATTACHED COMMENTS.

5525-83

Camranna's Note
Poor Copy

SCOTT'S GRASS KILLER because its granular consistency will give longer residual action and will effectively control resistant Chinch bugs, Sod Webworm (Lawn Moths), Ants and Ticks which infest lawns.

Damage may be prevented to your lawn if an application is put down in late Spring or early Summer before insects are visible.

FOR CHINCH BUGS:

Apply contents of this bag to 2500 square feet of lawn area. To insure best possible results, mow lawn, remove excess thatch and dead clippings, then water thoroughly before making application. Drench with water again after application so that granules will get down to the base of green plants. This is the area where Chinch Bugs will be found. Apply when insects appear and repeat in 7 to 10 days if necessary. It must be remembered that Chinch Bugs have several broods each summer so that repeat treatments at a later date may be necessary. The Chinch Bug is easily found on the very edge of the browned-out areas in the live grass at the base of the plant. They range in size from a pencil point to the adult bug which is about the size of a small ant and usually has a white spot on its back. (See picture on the front of the bag).

FOR SOD WEBWORMS (LAWN MOTHS), ANTS & TICKS:

Apply contents of this bag to 2500 square feet of lawn area. A second application 7 to 10 days later may be necessary.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Do not take internally. Avoid breathing dust. Harmful if inhaled or swallowed. Avoid contact with skin and eyes. For safety, wear gloves when applying. Wash thoroughly after using, particularly before smoking or eating. For eyes, flush profusely with water. If taken internally, induce vomiting. In both instances contact doctor. Keep pets and children off treated area until material is thoroughly washed in and completely dry. Avoid contamination of feed and food stuffs. Do not reuse container. Destroy when empty.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.

NOTE FOR PHYSICIAN:

Carbaryl is a moderate, reversible cholinesterase inhibitor.

NOTICE:

Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

SPREADER SETTINGS:

SCOTTS	7 1/2
LAWN BEAUTY	7 1/2
CYCLONE	3 3/4

NOTE: Above spreader settings are approximate. Allowances must be made for condition of spreader, humidity, speed at which applied, and roughness of area treated.

