

**GRO-WELL TOMATO & VEGETABLE DUST** with SEVIN\* and ZINEB is an insecticide-fungicide combination for the control of certain insects and diseases that infest tomatoes and certain vegetables.

**DIRECTIONS:** To dust, hold at 45° angle, lift hinged cap and squeeze uniformly. To avoid excessive dusting, do not point downward and make certain squeeze is uniform. This squeeze duster may be refilled from the 1 lb. or 46 oz. can of **GRO-WELL TOMATO & VEGETABLE DUST**.

Apply before insects become numerous and before diseases become apparent. Protect plants through the growing season by dusting at 7 to 10 day intervals and after heavy rains. Cover foliage thoroughly, but do not use in excess. Pay particular attention to cover underside of leaves. Use at the rate of 1 lb. per 1,200 sq. ft. (40' x 30') except on CUCUMBERS, SQUASH, MELONS and PEAS — use at the rate of 1 lb. per 2,000 sq. ft. (40' x 50').

VEGETABLES	INSECTS	DISEASES	DO NOT APPLY WITHIN —DAYS OF HARVEST
TOMATOES	Hornworm, Fruitworm, Leafhopper	Early & Late Blight, Anthracnose Gray & Septoria Leaf Spot	NO LIMITATION
PEPPERS	Flea Beetle & Tarnished Plant Bug	Anthracnose & Cercospora	NO LIMITATION
EGGPLANT		Phomopsis, Alternaria & Anthracnose	
CUCUMBER, SQUASH and MELON	Striped & Spotted Cucumber Beetle, Squash Bug, Flea Beetle & Leafhoppers	Downy Mildew, Anthracnose, Angular Leaf Spot and Scab	NO LIMITATION
BEANS	Mexican Bean, Flea, Bean Leaf & Japanese Beetles, Corn Earworm and Leafhoppers	Downy Mildew, Rust & Anthracnose	7 DAYS
BRUSSEL SPROUTS, CAULIFLOWER & BROCCOLI	Imported Cabbage Worm, Flea Beetle, Harlequin Bug, Corn Earworm, Armyworm & Young Cabbage Loopers (First or Second Instar Larvae)	Downy Mildew and Leaf Spot	7 DAYS
CABBAGE <sup>1</sup>			SEE NOTE 1
KOHLRABI <sup>2</sup>			SEE NOTE 2
LETTUCE		Botrytis Slime & Downy Mildew	SEE NOTE 3
CARROTS	Leafhoppers	Early & Late Blight	7 DAYS
PEAS	Pea Weevil	Anthracnose, Downy Mildew & Rust	10 DAYS
SWEET CORN	European Corn Borer <sup>4</sup> & Corn Earworm <sup>5</sup>		7 DAYS

**NOTES:** 1. Do not apply within 7 days of harvest. On Chinese Cabbage do not apply within 14 days of harvest. 2. Do not apply after edible parts are half grown. 3. Do not apply within 5 days of harvest on Head Lettuce or within 14 days on Leaf Lettuce. 4. Dust every 5 to 7 days from the time eggs are about to hatch. 5. Dust silks as soon as the silks appear, making at least three applications at 2 to 3 day intervals. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**CAUTION: KEEP OUT OF REACH OF CHILDREN & PETS. HARMFUL IF SWALLOWED OR INHALED.** May cause irritation of nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Wash hands thoroughly before eating or smoking. Do not contaminate feed and foodstuffs. Do not apply or allow to drift into areas occupied by unprotected persons or beneficial animals. If spilled, flush area with plenty of water. As this container can be refilled, we suggest that you use nothing but the **GRO-WELL TOMATO & VEGETABLE DUST**, as outlined above, or dispose of this container when empty.

Do not contaminate fish ponds. Do not apply where runoff is likely to occur. Do not use on crops to be fed to dairy animals or to animals being finished for slaughter.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**NOTE:** Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

\*Trade Mark of Union Carbide Corp. for 1-Naphthyl-N-methylcarbamate.

NET WEIGHT: 10 OUNCES  
SOLD BY: \_\_\_\_\_ EPA Reg. No. 5535-85

**ACCEPTED**

*Mar. 23, 1972*

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT

C 346 RLD

BLACK GREEN )



*Squeeze-Duster*


SINCE 1855

FOR TOMATOES & CERTAIN VEGETABLES

CONTAINS:

SEVIN\* and ZINEB

INSECTICIDE—FUNGICIDE



**ACTIVE INGREDIENTS:**

\*Carbaryl (1-Naphthyl N-methylcarbamate) ..... 5.0%  
Zineb (Zinc ethylene bisdithiocarbamate) ..... 4.8%  
(metallic Zinc content — .92%)

INERT INGREDIENTS ..... 90.20%

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

SEE BACK PANEL FOR ADDITIONAL REQUIRED PRECAUTIONS