

75-26 Pm-2 1 of 2
ACCEPTED

JUL 11 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 5535-96

Since 1855

GRO-WELL.

TOMATO AND VEGETABLE DUST

Controls Insects & Diseases Attacking Cucumbers, Eggplant, Peppers, Tomatoes, Leafy-Type and other Vegetables †

Makes Over 25 Gallons of Spray

ACTIVE INGREDIENTS:

* Carbaryl (1-naphthyl N-methylcarbamate)	5.00%
Zineb (Zinc ethylene bisdithiocarbamate) (metallic Zinc content - .92%)	4.50%

INERT INGREDIENTS:	90.50%
TOTAL	100.00%

* Registered trademark of Union Carbide Corp. for 1-naphthyl-N-methylcarbamate

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY INFORMATION**

NET WEIGHT
46 OZ (2 LB 14 OZ)

† As Listed

SOLD BY:
J & L ADKES, INC.
JAMAICA, N.Y. 11423

FRONT PANEL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: HARMFUL IF SWALLOWED OR INHALED. May cause irritation of nose, throat and skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED** - Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Never give anything by mouth to an unconscious or convulsing person.
- IF ON SKIN** - Wash with soap and water. Get medical attention if irritation persists.
- IF IN EYES** - Flush with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees, exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using. Use strictly in accordance with label cautions, warnings and directions. Apply this product only as specified on this label.

STORAGE: Store in original can in a locked storage area, away from other pesticides, fertilizer, food, and feed. Keep can tightly closed after opening. If broken or spilled, sweep up and discard in trash. Do not store diluted spray mixture. Make new dilution for each use.

PESTICIDE DISPOSAL: Securely wrap original can in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse can. Rinse thoroughly before discarding in trash.

GRO-WELL TOMATO & VEGETABLE DUST with SEVIN & ZINEB is an insecticide-fungicide combination for the control of certain insects and diseases that infest TOMATOES and certain VEGETABLES

Apply before insects become numerous and before diseases become apparent. Protect plants through the growing season by dusting at 7-10 day intervals and after heavy rains. Cover foliage thoroughly but do not use in excess. Pay particular attention to cover underside of leaves.

DUSTING: Use at the rate of 1 lb. per 1,200 sq. ft. (30' x 40') except on CUCUMBERS, MELONS, PEAS and SQUASH - use at the rate of 1 lb. per 2,000 sq. ft. (40' x 50')

SPRAYING: Add required material to water and stir thoroughly. Pour mixture into sprayer and agitate while spraying in order to obtain a thorough and even coverage. A sprayer which attaches to a hose, such as the GRO-WELL SHRUB & TREE SPRAYER, may be used. Use diluted spray at the rate of 1 gallon per 400 sq. ft. (20' x 20') for most pests listed - for pests marked ‡ use at the rate of 1 gallon per 200 sq. ft. (10' x 20').

DILUTION: 6-7 LEVEL TABLESPOONFUL PER GALLON OF WATER (1 POUND PER 10 GALLONS OF WATER)

FOR USE ON	SEE NOTE	INSECTS CONTROLLED	DISEASES CONTROLLED
Tomatoes	6	Fruitworm; Hornworm; Leafhopper	Anthraxosis (Blight) (Early & Late); Leaf Spots (Gray & Septoria)
Eggplant			Alternaria; Anthracnose; Phomopsis
Peppers	8	Flies Beets; Tarnished Plant Bug ‡	Anthraxosis; Cercospora
Cucumber Melon Squash	6	Beetles ‡ (Flea, Spotted & Striped Cucumber); Leafhopper ‡; Squash Bug ‡	Angular Leaf Spot; Anthracnose; Downy Mildew; Scab
Beans		Beetles ‡ (Bean Leaf); Flea ‡; Japanese ‡ & Mexican Bean ‡; Corn Earworm ‡; Leafhopper ‡	Anthraxosis; Downy Mildew; Rust
Broccoli Brussels Sprouts Cauliflower	1	Armyworm ‡; Corn Earworm ‡; Flea Beetle; Harlequin Bug; Impaired Cabbageworm ‡; Young Cabbage Looper ‡ (First or Second Instar Larvae)	Downy Mildew; Leaf Spot
Cabbage			
Kohlrabi	2		
Lettuce	3		Bacterial Stems; Downy Mildew
Carrots	1	Leafhoppers	Blight (Early & Late)
Peas	7		Anthraxosis; Downy Mildew; Rust
Sweet Corn	1-4.5	Corn Earworm ‡; European Corn Borer ‡	

WARRANTY STATEMENT

Follow directions carefully as to use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given hereon are not strictly adhered to. Seller gives no other or further warranty expressed or implied as to the merchantability or any other aspect of products sold.

NOTES: (1) Do not apply within 7 days of harvest. (2) Do not apply after seedling parts are well grown. (3) Do not apply within 10 days of harvest on head Lettuce or within 14 days on Leaf Lettuce. (4) Dust every 5-7 days from the time eggs are about to hatch. Apply thoroughly into the whorls when plants are about knee-high. Repeat at weekly intervals until tassels emerge. (5) Dust till as soon as the silks appear, making at least three applications at 2-3 day intervals. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. Do not feed treated forage to dairy animals or animals being finished for slaughter. (6) Do not apply within 5 days of harvest. (7) Do not apply within 10 days of harvest. (8) No pre-harvest treatments.