# GRO-WELL

1.19

# DANDELION AND BROADLEAF

WEED KILLER

4M-6-72

COVERS 2,000 sq. ft

KILLS
KNOTWEED, DANDELION, SPURGE,
CHICKWEED,
WILD ONION, POISON IVY AND
CERTAIN OTHER WEEDS

**CAUTION:**
KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

GRO-WELL DANDELION & BROADLEAF WEED KILLER may be used on BLUEGRASS, BEECGRASS, FESCUE, BERMUDA GRASS, ZOYSIA lawns, ocean lots, playgrounds and other similar areas where hard to kill weeds such as KNOTWEED, CHICKWEED, SPOTTED SPURGE, HENBIT, OXALIS, BLACK MEDIC and SHEEP SORREL are present in addition most other broadleaf weeds such as DANDELION and PLANTAIN and a partial list of other weeds listed below will be controlled.

**WHEN TO USE** Spray this lawn during the growing season when weeds are actively growing. Treat Wild Onion or Wild Garlic, CHICKWEED and HENBIT in the fall or in the spring as needed. On NEW LAWNS wait until the grass has reached a height of 2 inches before making an initial spray treatment. Wait 3 weeks before seeding new grass seed in bare spots. Poor lawn control may result if applied during drought or just before rain for best results. Make sure lawn is in good growing condition and mow at least one day before treatment. Do not water lawn within 24 hours after treatment.

**HOW MUCH TO USE** Mix 4 ounces of this mixture with 4 gallons of water and apply to 1,000 sq. ft. (20 x 50) of lawn area. A sprayer which attaches to a hose such as the GRO-WELL LAWN SPRAYER may be used for this purpose.

**WEEDS CONTROLLED**

| | | | | |
|---|---|---|---|---|
| BEGGSTRAW | DANDELION | LAMB'S-QUARTER | PLANTAINS | SPURGE |
| BLACK MEDIC | DOCK | LESPEDEZA | POISON IVY | SUMPWEED |
| BUCKHORN | GROUND IVY | MALLOW | POISON OAK | WILD CARROT |
| CHICORY | HOUSELEEK | MORNING GLORY | PURSLANE | WILD GARLIC |
| CHICKWEED | HENBIT | PENNYCRESS | RAGWEED | WILD LETTUCE |
| CLOVER | KNOTWEED | PIGWEED | SHEPHERD'S PURSE | WILD ONION |

YARROW and many others

**CAUTION** Harmful if swallowed. Keep out of reach of children and pets. Keep children and pets off treated area until completely dry. Avoid skin, eye and clothing contact. In case of contact flush with water. Avoid contamination of food or foodstuffs.

Do not allow spray or mist to get on vegetables, ornamental plants, flowers, shrubs, trees or other desirable plants. A coarse type spray is less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not pour spray solution near these plants. Do not spray on Carpetgrass, Dichondra or on lawns where desirable Bentgrass or Clovers are present. This product may be injurious to Centipede or St. Augustine grass lawns.

Do not contaminate domestic or irrigation water. Do not store near fertilizers, seeds, insecticides or fungicides. When this container is empty, wash thoroughly and destroy. Do not re-use for any other purpose.

**NOTE:** Follow directions carefully as use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given herein are not strictly adhered to.

**ACTIVE INGREDIENTS:**

| | |
|---|---|
| *Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid | 3.66% |
| **Dimethylamine salt of 2,4-dichlorophenoxyacetic acid | 8.07% |
| ***Dimethylamine salt of Beacon (3,6-dichloro-o-anisic acid) | 8.84% |
| Dimethylamine salts of related compounds | 8.11% |
| **INERT INGREDIENTS** | 67.32% |
| Total | 100.00% |

*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 3.02% or 0.26 lbs. per gallon
**2,4-dichlorophenoxyacetic acid equivalent 6.78% or 0.576 lbs. per gallon
***3,6-dichloro-o-anisic acid equivalent 8.65% or 0.06 lbs. per gallon

Sold by: NET CONTENT: 8 OUNCES EPA REG. NO. 5535-67

J. & L. ADIKES, INC., JAMAICA, NEW YORK 11423

ACCEPTED
Jan. 26, 1973
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE, AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER EPA REG. NO. 5535-67 SUBJECT TO ATTACHED COMMENTS.

5535-67/36

4M-6-72

# GRO-WELL

1.19

## DANDELION AND BROADLEAF WEED KILLER



COVERS 2,000 sq. ft

KILLS
KNOTWEED, DANDELION, SPURGE, CHICKWEED, WILD ONION, POISON IVY AND CERTAIN OTHER WEEDS

**CAUTION:**
KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

GRO-WELL DANDELION & BROADLEAF WEED KILLER may be used on BLUEGRASS, RYEGRASS, FESCUE, BERMUDA and ZOYSIA lawns, vacant lots, playgrounds, and other similar areas where hard to kill weeds such as KNOTWEED, CHICKWEED, SPOTTED SPURGE, HENBIT, OXALIS, BLACK MEDIC and SHEEP-SORREL are present in abscon, most other broadleaf weeds such as DANDELION and PLANTAIN, and a partial list of other weeds, listed below will be controlled.

WHEN TO USE: Spray any time during the growing season, when weeds are actively growing. Treat WILD ONION, WILD GARLIC, CHICKWEED and HENBIT in the fall or in the spring, as needed. ON NEW LAWNS wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before seeding new grass seed in bare spots. Poor weed control may result if applied during drought or rest before rain. For best results make sure lawn is in good growing condition and mow at least one day before treatment. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE: Use 4 ounces of this material with 4 gallons of water and apply to 1,000 sq. ft. (20' x 50') of lawn area. A sprayer which attaches to a hose, such as the GRO WELL LAWN SPRAYER, may be used for this purpose.

WEEDS CONTROLLED

| | | | | |
|---|---|---|---|---|
| BEDSTRAW | DANDELION | LAMBSQUARTER | PLANTAINS | SPEEDWELL |
| BLACK MEDIC | DOCK | LESPEDEZA | POISON IVY | SPURGE |
| BUCKHORN | GROUND IVY | MALLOW | POISON OAK | WILD CARROT |
| CHICORY | HENBIT | MORNING GLORY | PURSLANE | WILD GARLIC |
| CHICKWEED | KNOTWEED | PEPPERGRASS | RAGWEED | WILD LETTUCE |
| CLOVER | | PURSLANE | SHEPHERD'S PURSE | WILD ONION |

YARROW and certain others

CAUTION: Harmful if swallowed. Keep out of reach of children and animals. Keep children and pets off treated area until completely dry. Avoid skin, eye and clothing contact. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

Do not allow spray or mist to get on vegetables, ornamental plants, flowers, shrubs, trees or other desirable plants. A coarse type spray is less likely to drift. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not pour spray solution near these plants. Do not spray on Carpetgrass, Dichondra or on lawns where desirable Bentgrass or Clovers are present. This product may be injurious to Centipede or St. Augustine grass lawns.

Do not contaminate domestic or irrigation water. Do not store near fertilizers, seeds, insecticides or fungicides. When this container is empty, wash thoroughly and destroy. Do not re-use for any other purpose.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given hereunto are not strictly adhered to.

ACTIVE INGREDIENTS:

| | |
|---|---|
| *Dimethylamine salt of 2 (2 methyl 4 chlorophenoxy) propionic acid | 3.68% |
| **Dimethylamine salt of 2 4 d chlorophenoxyacetic acid | 2.07% |
| ***Dimethylamine salt of Dicamba (3 6-dichloro o anisic acid) | 0.66% |
| Dimethylamine salts of related components | 0.11% |
| INERT INGREDIENTS | 93.32% |
| Total | 100.00% |

*2 (2-methyl 4 chlorophenoxy) propionic acid equivalent 3.02% or 0.26 lbs per gallon
**2,4 d chlorophenoxyacetic acid equivalent 6.70% or 0.576 lbs. per gallon
***3 6 dichloro o anisic acid equivalent 0.65% or 0.05 lbs per gallon

Sold by: NET CONTENT: 8 OUNCES EPA REG. NO. 5535-87

J. & L. ADIKES, INC., JAMAICA, NEW YORK 11423

ACCEPTED
Jan. 26, 1973
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 5535-87 SUBJECT TO ATTACHED COMMENTS.

5535-27/86

5