


GRO-WELL

DANDELION AND BROADLEAF WEED KILLER

27
35
KILLS CHICKWEED, CLOVER
KNOTWEED, POISON IVY, SPEEDWELL,
WILD ONION
AND OTHER LAWN WEEDS *

TREATS
4,000 SQ. FT. (50' X 80')



ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-(2-methyl-4-chlororhexoxy)triazine	3.66%
**Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	8.07%
***Dimethylamine salt of picloric (3,6-dichloro-o-anisic acid)	0.84%
Dimethylamine salts of related compounds	0.11%

INERT INGREDIENTS: 87.32%

TOTAL 100.00%

*2,4-(2-methyl-4-chlororhexoxy)triazine acid equivalent 0.02% or 0.26 lbs. per gallon
 **2,4-dichlorophenoxyacetic acid equivalent 0.70% or 0.576 lbs. per gallon.
 ***3,6-dichloro-o-anisic acid equivalent 0.09% or 0.09 lbs. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY INFORMATION

NET CONTENTS

16 Fl.oz. (1 Pt.)

as listed

SINCE 1855

SOLD BY: J. & L. ADIKES, INC.
JAMAICA, N.Y. 11423

11/3

REAR PANEL

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS

TYPE SIZE

----- PRECAUTIONARY STATEMENTS

----- HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. KEEP OUT OF REACH OF CHILDREN and pets, and keep them off treated areas until completely dried. HARMFUL IF SWALLOWED. In case of contact with eyes, flush profusely with water, get medical attention if irritation persists. If swallowed, drink one or two glasses of water. Induce vomiting by placing finger in back of throat. Call a physician. Never give anything by mouth to an unconscious person. Avoid breathing of spray mist and contact with skin, eyes and clothing. Wash off immediately with soap and water. ✓

Read entire label before using. Use strictly in accordance with label cautions and directions. Apply this product only as specified on this label.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water used for irrigation or domestic purposes. ✓

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

-----STORAGE AND DISPOSAL

STORAGE: Store in original bottle in a locked storage area, away from other pesticides, fertilizer, food, and feed. Keep bottle tightly closed after opening. If broken or spilled, apply an absorbent, sweep up and discard in trash. Do not store diluted spray material, make a new dilution for each use.

PESTICIDE DISPOSAL: Securely wrap original bottle in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

----- WARRANTY STATEMENT


Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given herewith are not strictly adhered to. Seller gives no other or further Warranty, expressed or implied as to the merchantability or any other aspect of products sold.

PRODUCT:

MANUFACTURED BY: ...

GRO-WELL® DANDELION & BROADLEAF WEED KILLER may be used on LAWNS, VACANT LOTS, PLAYGROUNDS, and other similar areas where hard-to-kill weeds are present. A sprayer which attaches to a hose, such as the GRO-WELL® HOSE SPRAYER, may be used with this product to control the partial list of weeds shown.

DOSAGE LISTED TO BE MIXED WITH 4 GALLONS OF WATER & APPLIED TO 1,000 SQ. FT. (20'x50')

FOR USE ON	SEE NOTE	TO CONTROL			DOSAGE (Application Rate)	
<p>✓ (contains Bluegrass, Ryegrass, Bermuda, Fescue)</p>	1	Bedstraw	Henbit	Poison Oak	8 tbsp. (4 ozs.)	
		Black Medic	Knotweed	Purslane		
		Buckhorn	Lamb's-quarter	Ragweed		
		Chicory	Lamb's-ear	Shepherd's Purse		
		Chickweed	Mallow	Speedwell		
		Clover	Nutmeg	Sourgrass		
		Dandelion	Pigweed	Wild Carrot		
		Dock	Plantain	Wild Garlic		
		Ground Ivy	Poison Ivy	Wild Lettuce		
		Heal-all		Wild Onion		
				Yarrow and certain other weeds		
						

NOTE: (1) spray any time during the growing season, when weeds are actively growing. Treat Wild Onion, Wild Garlic, Chickweed and Henbit in the Fall or in the Spring, as needed. On NEW LAWNS wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 2 weeks before putting new grass seed in bare spots. Poor weed control may result if applied during drought or just before rain. For best results, make sure lawn is in good growing condition and mow at least one day before treatment. Do not water lawn within 24 hours after treatment.

CAUTION: do not contaminate domestic or irrigation water. Do not allow spray or mist to get on vegetables, ornamental plants, flowers, shrubs, trees or other desirable plants. A coarse spray is less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not pour spray solution near these plants. Do not spray on Carpetgrass, Dichondra or on lawns where desirable Bentgrass or Clovers are present. This product may be injurious to Centinoda and St. Augustinegrass lawns.

EPA Reg.No. 5035-r

EPA Est. 72-PA-1

50/3