

UP
5,000 SQ. FT.

Controlled
TUPERAN

YOU CAN SEED WITH APPLICATION

OTUP... Co., Inc. ... KILLER

... OUT ...
... OR AD...

NET WEIGHT 10 LBS

EPA Est. No. 1159

EPA Reg. No. 553

ADIKO, INC. MAICA N.Y. 114

GRO-WELL

PRE-VENT. CRABGRASS CONTROL

GRO-WELL® PRE-VENT® CRABGRASS CONTROL may be applied in conjunction with seeding if directions given below are followed.

GRO-WELL PRE-VENT will effectively and selectively control CRABGRASS (Smooth & Hairy) as well as other annual weeds such as FOXTAIL and BARNYARD GRASS. PRE-VENT may be used on BLUEGRASSES, FESCUES, BENTGRASSES (Penncross, Seaside & Astoria), and PERENNIAL RYEGRASSES without danger of injury when applied as directed. Do not use on other Bentgrass strains, nor on warm season grasses such as Bermuda Grass as injury might occur. PRE-VENT should be applied in the Spring before Crabgrass and other annual weeds start to grow, following spreading directions given below.

HOW TO APPLY: Prior to application, lawn should be raked to remove extraneous material and will therefore insure uniform application. In order to obtain expected results PRE-VENT should be evenly applied with no skips or misses following spreading instructions listed below.
NOTE: Do not rake lawn or disturb soil surface after application.

FOR USE WITH SEEDING: Apply contents of this bag to 5,000 sq. ft. (50' x 100') of lawn area, using spreader setting given below WITH SEEDING. Apply immediately after seeding, before crabgrass and other annual weeds germinate. For best results repeat application in four weeks. Make certain to provide sufficient moisture to insure germination of new grass seed. Follow watering directions carefully.

FOR USE WITHOUT SEEDING: Apply contents of this bag to 2,500 sq. ft. (50' x 50') of lawn area, using spreader setting given below WITHOUT SEEDING. Follow watering directions carefully.

non-carcinogenic.

USE OF FERTILIZER: A normal application of fertilizer may be used either before or after application of PRE-VENT, following directions listed on the label. This will encourage the growth of the grass enabling it to fill in the bare areas more rapidly.

SPREADING INSTRUCTIONS: It is recommended that a cross-pattern be used, (i.e., once in a north/south direction, then immediately followed by an east/west direction) using the spreader settings listed below, continuing until proper amount has been applied. This method of application will give a more uniform coverage for maximum control. If setting for your particular spreader is not listed below, open spreader approximately 1/3 so that material flows out uniformly, but does not pour out.

SPREADER SETTINGS

	APPLY CONTENTS TO	CYCLONE	SCOTTS
WITH SEEDING	5,000 sq. ft.	5	8
WITHOUT SEEDING	2,500 sq. ft.	8	12

NOTE: Variations in equipment & material, as well as accuracy of machine adjustment, may cause actual application rates to vary slightly. These suggested settings are for your guidance only, therefore, we make no warranty, expressed or implied as to the uniformity of coverage obtained from settings listed above.

WATERING DIRECTIONS: After application, water area sufficiently to wash PRE-VENT off grass blades to carry it into the root zone of germinating weeds, being careful not to apply too much water as puddles might form and chemical will float away. For best results, area should get a 1/2" of water, either by rainfall or irrigation within 3 days after application.

CAUTION: KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. Keep children and pets off treated area until it is completely dried. Harmful if inhaled. Avoid breathing dust. May irritate eyes, nose, throat and skin. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. If spilled, immediately sweep area and dispose of sweepings. Do not reuse container. Destroy when empty, either by burying with waste, away from domestic water supplies, or by burning. Stay away from smoke and fumes.

Apply only to lawn areas and not to grass areas which might be used for grazing or harvesting for feed and crops. Do not apply to areas containing flowers, shrubs, vegetables and fruits. When mowing lawn for first time after application, carefully dispose of all clippings collected.

ENVIRONMENTAL CAUTIONS:

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply to areas where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

REGAL

Supply & Chemical
P. O. Box 1055, El Paso, Texas

NOTICE: FOR INDUSTRIAL, INSTITUTIONAL OR PROFESSIONAL USE ONLY

● WATER SOLUBLE

● HIGH CONCENTRATE

ENDWEED

DIRECTIONS FOR USE:

Uniformly spray one gallon of product mixed in 10 gals. of water over 1000 sq. ft. of soil surface. Thoroughly wet all foliage to insure contact action. Best results are obtained when applications are made during or soon after weed emergence while growth is succulent.

Where vegetation is tall and dense, use 1 1/2 gals. of product in 10 gals. of water per 1000 sq. ft. For control of hard-to-kill perennial weeds and grasses such as Johnson Grass, Bindweed and Wild Carrot, use 3 to 4 1/2 gals. per 1000 sq. ft. Although this product has considerable activity through foliar contact, much of its action is through the roots and therefore, adequate rainfall or watering to move the chemical into the root zone is necessary.

Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed-free soil surface.

Care must be taken to keep this herbicide away from desirable flower beds, lawns and shrubs. Do not apply over roots of valuable trees, or woody ornamental plants. Only apply herbicide outside of the drip line of trees. Avoid drift to desirable plants. Do not use the same spray equipment for other purposes.

RESIDUAL ACTION
VEGETATION KILLER

PER SITE

FOR NON-SELECTIVE WEED CONTROL ON NON-CROP-LAND AREAS SUCH AS PARKING LOTS, FENCE ROWS, RAILROAD SIDINGS, STORAGE YARDS, AROUND BUILDINGS OR INDUSTRIAL SITES.

ACTIVE INGREDIENTS:

Prometon [2,4-bis(isopropylamino)-6-methoxy-s-triazine]*	3.76%
Pentachlorophenol	1.76%
2,3,4,6-Tetrachlorophenol	0.24%
Petroleum distillates	78.92%
INERT INGREDIENTS:	15.32%

*Pramitol[®], trademark of CIBA-GEIGY Corp.

Good industrial safety practice should include the use of rubber gloves, protective clothing and face mask or eye shield when using this product or any other chemical.



SAFETY PRECAUTIONS ON LABEL MUST BE FOLLOWED TO MEET OSHA STANDARDS.

EPA Reg. No. 5576-31
EPA Est. No. 5576-Tx-1

IMPORTANT

Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Since weather and other conditions under which this product may be used will vary, the seller makes no warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risks of use or handling not in accordance with directions or suggestions.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

- Harmful if swallowed. Avoid contact with eyes and skin.
- In case of skin contact wash with soap & water.
- Avoid inhalation of spray mist. Avoid contamination of feed and food.
- Do not spill, pour or store near heat or open flame.
- Do not contaminate water used for irrigation or domestic purposes.
- Do not reuse container. Destroy when empty.
- Store at temperatures above 32°F.

FLAMMABLE MIXTURE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NET CONTENTS _____

Jan. 13, 1975

5576-31

FORMULATED FOR INDUSTRIAL AND INSTITUTIONAL USE

STRILATE

FOR USE IN AND AROUND ASPHALT
BARE GROUND WEED CONTROL

DIRECTIONS FOR USE:

Uniformly spray 2 1/2 - 3 gals. per 1000 sq. ft. of soil surface, undiluted (100-120 gals. per acre.) Thoroughly wet all foliage to insure contact action. Best results are obtained when applications are made during or soon after weed emergence while growth is succulent. Where vegetation is tall and dense, use 5 gals. per 1000 sq. ft. (200 gals. per acre.) For control of hard-to-kill perennial weeds and grasses such as Johnson Grass, Bindweed and Wild Carrot, use 8-12 gals. per 1000 sq. ft. Although this formulation has considerable activity through foliage contact, much of its activity is through the roots and therefore adequate rainfall or watering to move the chemical into the root zone is necessary. Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed free soil surface.

STRILATE may be mixed with standard and special cutback asphalt, (such as RC, MC and SC.) STRILATE should be used at the rate of 9-13 1/2 fl. oz. per sq. yd. STRILATE should be simultaneously mixed with the cutback asphalt as it is being pumped into the applicator or transport truck to assure a homogeneous mix. Temperatures in the applicator or transport truck should not exceed 170° F. STRILATE may be applied under asphalt to extend the life of such coatings. STRILATE should be used only where the area to be treated has been prepared according to good construction practices. STRILATE will not control woody vegetation such as small trees, bush or woody vines.

IMPORTANT

Read the entire Directions for use and the conditions of sale and warranty before using this product. Since weather and other conditions under which this product may be used will vary, the seller makes no warranty of any kind, express or implied, concerning the use of this product. The user assumes all risks of usage or handling not in accordance with directions or suggestions.


EPA Reg. No. 5576-32 EPA Est. No. 5576 Tx-1

NET CONTENTS _____


ACTIVE INGREDIENTS:

*Prometon (2,4-bis (isopropylamino)-6-methoxy-s-triazine)	1.5%
Pentachlorophenol	0.7%
2,3,4 6-Tetrachlorophenol	0.1%
Petroleum distillate	93.2%
INERT INGREDIENTS	4.5%

*Prometon® , trademark of CIBA-GEIGY Corporation



WITHOUT "STRILATE"



WITH "STRILATE"

ACCEPTED
12-10-74
5576-32
Comments

UNDER THE FEDERAL INSECTICIDE ACT
REGISTERED

REGAL

Supply & Chemical
Box 1905 El Paso, Texas

AROUND:

- GUARD RAIL
- MACHINERY
- STORAGE AC
- MILITARY
- HIGHWAY ME
- INDUSTRIAL
- DRAINAGE D

GENERAL INFO

THIS PRODUCT IS A SPECIFIC KILLER FOR CONTROLLING WEEDS AND GRASSES. APPLICATED IN AREAS WHERE STERILE BARE GROUND WITH THE AMOUNT OF CONTACT WITH OTHER CONDITIONS. THE PRODUCT WILL CONTROL WEEDS ON ALL SURFACES. ESPECIALLY COMBINATION WEED CONTROL.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. CONTAINS PETROLEUM PRODUCTS WHICH ARE HARMFUL OR FATAL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES, OR CLOTHING. IF CONTACTED WITH SKIN, WASH WITH WATER AND SOAP. IF CONTACTED WITH EYES, FLUSH WITH WATER FOR SEVERAL MINUTES. CONSULT A PHYSICIAN. AVOID INHALATION OF VAPORS. AVOID CONTAMINATION OF FOOD AND FEED. DO NOT REUSE CONTAINERS FOR OTHER PURPOSES. DO NOT REUSE CONTAINERS FOR OTHER PURPOSES. DO NOT USE, POUR, OR DISPOSE OF IN WATER. STORE AT TEMPERATURES BELOW 100° F. HARMFUL IF SWALLOWED.

This product is toxic to fish and aquatic life. Do not apply to ponds. Do not apply to water. Do not apply to water by cleaning. Apply this product only to bare ground. Good industrial safety practices require protective clothing and mask when using other chemical.