

PRE-VENT

WEED CONTROL



PREVENTS GROWTH OF CERTAIN WEEDS AROUND

VEGETABLES, FLOWERS & SHRUBS



CRABGRASS



CHICKWEED



SPURGE

ACTIVE INGREDIENT:

*Dimethyl Tetrachloroterephthalate..... 6.00%

INERT INGREDIENTS: 94.00%

TOTAL 100.00%

* DACTHAL is a Trade Mark of Diamond Shamrock Chemical Co.
 * Use as a pre-emergent herbicide patented under U.S. Patent No. 2,923,634.

CAUTION: KEEP OUT OF REACH OF CHILDREN
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET WT : 16 OZ. (1 LB.)

If cultivation is desired for appearance make or to break crusted soil to allow penetration of moisture, it is recommended that a second application be made at that time. If rain does not occur within 3 to 4 days after application a light watering of the area will increase the effectiveness.

FOR USE AROUND VEGETABLES: This product may be used around the following vegetables, making certain that you apply only as specified, carefully following the instructions and precautions as noted.

VEGETABLES	WHEN TO APPLY	SEE NOTE
BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE	AT SEEDING OR TRANSPLANTING	1
GARLIC, ONIONS		2, 3
SEEDED MELONS (Cantaloupe, Honeydew, Watermelons), CUCUMBERS, SQUASH (Summer and Winter)	ONLY WHEN PLANTS HAVE 4 TO 5 TRUE LEAVES	4 - 5 - 6
COLLARDS, KALE, MUSTARD GREENS, TURNIPS (Greens and Roots)	AT SEEDING	1
LETTUCE (Leaf)	1-3 WEEKS AFTER EMERGENCE	3
LETTUCE (Head)	1-6 WEEKS AFTER SEEDING	3
TOMATOES, EGGPLANTS, PEPPERS	4-6 WEEKS AFTER TRANSPLANTING	4, 7
STRAWBERRIES (New Plantings)	AT TRANSPLANTING	1, 6

Notes: 1. Material can be applied prior to planting. 2. A side application can be made on Onions only up to 14 weeks after planting or transplanting. 3. Preplant application not recommended. 4. Plants should be well established and growing conditions favorable. 5. If applied earlier than recommended and/or growing conditions are unfavorable, crop injury may result. 6. Soil incorporation not recommended. 7. Do not apply to seeded plants until they are 4" to 6" high, well established, and growing conditions are favorable. 8. Do not apply while plants are fruiting.

MAY BE USED IN BEDS CONTAINING THE FOLLOWING SHRUBS, FLOWERS, TREES & ORNAMENTALS: For best results, apply in early Spring after plants have become established. No waiting period is required before applying to transplanted shrubs or bushes.

- | | | | |
|-----------------|--------------------|-------------------|---------------------|
| ABELIA | CYPRESS | LANTANA | REDBUD |
| AGERATUM | DAHLIA | LAVENDER COTTON | RHODOENDRON |
| AGUGA | DELPHINIUM | LILAC | ROSES |
| ALYSSUM | DEUTZIA | LILY | ROSE MOSS |
| ARBORVITAE | DOGWOOD | LOCUST | RUSSIAN OLIVE |
| ASH | ELM | LUPINE | SALVIA |
| ASTER | ENGLISH IVY | MAGNOLIA | SCARLET SAGE |
| AZALEA | EUONYMUS | MAPLE | SNAPDRAGON |
| BABYSBREATH | EVENING PRIMROSE | MARIGOLD | SPIREA |
| BARBERY | FAILLARDIA | MOCK ORANGE | SPRUCE |
| BELLFLOWER | FERNLEAF YARROW | MORNING GLORY | STONE CROP |
| BIRCH | FEVERFEW | MOTHER OF THYME | STRAWFLOWER |
| BLEEDING HEART | FIR | MOUNTAIN LAUREL | SUNDRIPS |
| BLOOD LEAF | FORGET ME NOT | MOURNING BRIDE | SUNFLOWER |
| BOXWOOD | FORSYTHIA | MOURNING PINKS | SWEET PEA |
| BUGLOSS | FOUR O'CLOCK | NASTURTIUM | SYCAMORE |
| CAMELLIA | FOXGLOVE | OAK | TREE PEONY |
| CANDLE LARKSPUR | GERANIUM | ORFANE | TULIP TREE |
| CANDYTUFTS | GLADIOLUS | PACHISTIMA | VARIEGATED PRIVET |
| CHESTNUT | GOLD DUST | PACHYSANDRA | VERBENA |
| CHRYSANTHEMUM | GOLDEN MARGUERITE | PEONIES | VIOLLET |
| CINQUEFOIL | GOLDEN TUFT | PETUNIA | VIBURNUM |
| COLEUS | GUM | PINE | VIRGINIA SPIDERWORT |
| COLUMBINE | HAWTHORN | PINK HEATH | WALNUT |
| CORAL BELLS | HOLLY | PITTOSPORUM | WEIGELA |
| COREOPSIS | HONEYSUCKLE | PODDCARPUS | WILLOW |
| COSMOS | HYDRANGEA | POKER PLANT | WILSON VIV |
| COTONEASTER | IRIS | POPLAR | WORMWOOD |
| COTTONWOOD | JAPANESE ANDROMEDA | PRIVET | YEW |
| CRABAPPLE | JUNIPER | PURPLE CONEFLOWER | ZINNIA |

SHOULD NOT BE USED ON THE FOLLOWING PLANTS: Bugle Weed, Button Pink, Carnation, Gum, Germander, Mesebryanthemum, Pansy, Philox, Sweet William and Tetanthera.

LIST OF WEEDS CONTROLLED

- | | | |
|------------------------------|---------------------------|------------------------|
| ANNUAL BLUEGRASS | GREEN FOXTAIL | PURSLANE |
| BROWN TOP PANICUM | JOHNSON GRASS (from Seed) | REDROOT PIGWEED |
| CARPETWEED | LAMBSQUARTER | SESBANIA (Microcarpal) |
| CHICKWEED (Common) | LOVEGRASS | TEXAS MILLET |
| CRABGRASS (Hairy and Smooth) | NODDING SPURGE | WITCHGRASS |
| FLORIDA PUSSLEY | | YELLOW FOXTAIL |

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking. Do not store near feed foodstuffs, seeds, fertilizers or other pesticides. Store in a dry place. Do not graze on treated areas or feed plant refuse to livestock. Do not reuse container. Destroy when empty, either burying with other wastes away from water supplies, or by burning. Do not inhale smoke or fumes.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given herewith are not strictly adhered to.

SOLD BY:

5555-102
 J. & L. ADIKES, INC.

EPA Reg. No. 5535-102

J. & L. ADIKES, INC. JAMAICA, NEW YORK 11423

ACCEPTED WITH COMMENTS

4M-11-75

EPA Est. No. 1159-NJ-1