


GRO-WELL. DANDELION BROADLEAF

CONTAINS *TRIMEC® HERBICIDE



SPECIALLY FORMULATED TO CONTROL
DANDELION, CHICKWEED, KNOTWEED
AND CERTAIN OTHER HARD-TO-KILL WEEDS

ACTIVE INGREDIENTS:	
* Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.178%
** Diethanolamine salt of 2,4-dichlorophenoxyacetic acid	0.177%
*** Diethanolamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.016%
INERT INGREDIENTS:	99.629%
TOTAL	100.000%

* Equivalent to 2-(2-methyl-4-chlorophenoxy) propionic acid 0.120%
 ** Equivalent to 2,4-dichlorophenoxyacetic acid 0.120%
 *** Equivalent to 3,6-dichloro-o-anisic acid 0.0114%

WARNING:

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL
FOR ADDITIONAL CAUTIONS

NET WEIGHT: 18 FL. OZ. (1 LB. 2 OZ.)

EPA Est. No. 11715-TN-1

SHAKE WELL BEFORE USING
READ ENTIRE LABEL BEFORE USING

Use GRO-WELL® DANDELION & BROADLEAF WEED KILLER SPRAY to spot-treat weeds in the lawn. For best results apply when weeds are actively growing. Do not water within 24 hours after spraying. Effects begin to show after a few days and weeds gradually die. Up to 4 weeks may be required for complete kill. Repeat in 3-4 weeks if necessary.

DIRECTIONS

Shake can well before using. This product is designed to permit easy treatment of lawn weeds without bending or stooping. Remove protective cap, aim at weed and depress valve, making certain valve opening is pointed away from you. Only a short burst is needed. Over spraying only wastes the spray.

WEEDS CONTROLLED		
BEDSTRAW	FILAREE	PENNYWORT
BUCKHORN	FLEABANE	PEPPERWEED
BLACK MEDIC (YELLOW CLOVER)	FLORIDA PUSLEY	PIGWEEED
CANADA THISTLE	GROUND IVY (CREEPING CHARLIE)	PLAINTAINS
CHICKWEED	HEAL-ALL	POISON IVY
CLOVER	HENBIT	PRICKLY LETTUCE
CUDWEED	KNOTWEED	PURSLANE
CURLED DOCK	LAMBSQUARTERS	RAGWEED
DANDELION	MATCHWEED	SHEEP SORREL
DOLLAR WEED	MUSTARD	SHEPHERD'S PURSE
ENGLISH DAISY	OXALIS	SPURGE
		YARROW

WARNING: KEEP OUT OF REACH OF CHILDREN. HARMFUL OR FATAL IF SWALLOWED. HARMFUL IF INHALED. Avoid contact with skin, eyes, or clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Avoid breathing of vapor or spray mist. If swallowed, do not induce vomiting; call physician immediately. Wash well after using and before eating or smoking.

Avoid spraying susceptible plants (flowers, shrubs, etc.). Bentgrass, Carpetgrass, and St. Augustine Grass may be injured if contacted. Do not apply within dripline of desirable plants. Avoid accidental drift or application to nearby garden plants, flowers, or shrubs. Do not store near seed, fertilizers, insecticides, or fungicides. Do not contaminate water used for irrigation or domestic purposes.

WARNING: Contents of can under pressure. Do not puncture. Do not use or store near heat or open flame. Exposures to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.


NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller; the buyer assumes all risks if directions given herewith are not strictly adhered to.

*TRIMEC® is a registered Trademark of P.B.I. - GORDON CORP. - U.S. PAT. NO. 3,284,186

Sold By:

E.P.A. Reg. No. 5535-98

J. & L. ADIKES, INC. JAMAICA, N. Y. 11423



10M-10-74

GRO-WELL®


PRE-VENT

WEED CONTROL

SINCE 1885

PREVENTS GROWTH OF CERTAIN WEEDS
AROUND

VEGETABLES, FLOWERS & SHRUBS



CRABGRASS

CHICKWEED

SPURGE

ACTIVE INGREDIENT:

*Dimethyl Tetrachloroterephthalate..... 6.00%

INERT INGREDIENTS:..... 94.00%

TOTAL 100.00%

* DACTHAL is a Trade Mark of Diamond Shamrock Chemical Co.
* Use as a pre-emergent herbicide patented under U.S. Patent No. 2,923,634.

CAUTION: KEEP OUT OF REACH OF CHILDREN

GRO-WELL® PRE-VENT WEED CONTROL is for use as a pre-emergent application for the control of CRABGRASS, COMMON CHICKWEED, ANNUAL GRASSES and other weeds listed below on mineral soils in VEGETABLE GARDENS and FLOWER & SHRUB BEDS.

DIRECTIONS: Punch out only two of the perforated marks on top of can. Apply contents of this can evenly by shaking over 250 sq. ft. (25' x 10') of area to be treated for the plants listed below, with the exception of STRAWBERRIES, which should be applied to a 10' x 33' or larger area. For a larger area, a second satisfactory result may be obtained.

This product will not kill existing weeds but will control weed seeds as they germinate and emerge from the soil. Before application, cultivate soil to provide a weed free and uniform surface. Application should be made within two to three days following cultivation, to prevent sub-surface germinating weeds from becoming established. One application should last for three months or more. A second application in late Summer or early Fall may be necessary for late germinating weeds. Do not disturb soil surface after application as this will reduce effectiveness. If cultivation is desired for appearance sake or to break crusted soil to allow penetration of moisture, it is recommended that a second application be made at that time. If rain does not occur within 3 to 4 days after application a light watering of the area will increase the effectiveness.

FOR USE AROUND VEGETABLES: This product may be used around the following vegetables, making certain that you apply only as specified, carefully following the instructions and precautions as noted.

VEGETABLES	WHEN TO APPLY	SEE NOTE
BROCCOLI, BRUSSELS SPROUTS CAULIFLOWER, CABBAGE	AT SEEDING OR TRANSPLANTING	1
GARLIC, ONIONS		2-3
SEEDED MELONS (Cantaloupe, Honeydew, Watermelons), CUCUMBERS, SQUASH (Summer and Winter)	ONLY WHEN PLANTS HAVE 4 TO 5 TRUE LEAVES	4-5-6
COLLARDS, KALE, MUSTARD GREENS, TURNIPS (Greens and Roots)	AT SEEDING	1
LETTUCE (Leaf)	13 WEEKS AFTER EMERGENCE	3
LETTUCE (Head)	16 WEEKS AFTER SEEDING	3
TOMATOES, EGGPLANTS, PEPPERS	4-6 WEEKS AFTER TRANSPLANTING	4-7
STRAWBERRIES New Plantings	AT TRANSPLANTING	1-8

Notes: 1. Material can be applied prior to planting. 2. A side application can be made on Onions only up to 14 weeks after planting or transplanting. 3. Preplant application not recommended. 4. Plants should be well established and growing conditions favorable. 5. If applied earlier than recommended and/or growing conditions are unfavorable, crop injury may result. 6. Soil incorporation not recommended. 7. Do not apply to seeded plants until they are 4" - 6" high, well established, and growing conditions are favorable. 8. Do not apply while plants are fruiting.

MAY BE USED IN BEDS CONTAINING THE FOLLOWING SHRUBS, FLOWERS, TREES & ORNAMENTALS: For best results, apply in early Spring after plants have become established. No waiting period is required before applying to transplanted shrubs or bushes.

- | | | | |
|-----------------|--------------------|-------------------|---------------------|
| ABELIA | CYPRESS | LANTANA | REDBUD |
| AGERATUM | DAHLIA | LAVENDER COTTON | RHODODENDRON |
| AGUGA | DELPHINIUM | LILAC | ROSES |
| ALYSSUM | DEUTZIA | LILY | ROSE MOSS |
| ARBORVITAE | DOGWOOD | LOCUST | RUSSIAN OLIVE |
| ASH | ELAEGNUS | LUPINE | SALVIA |
| ASTER | ELM | MAGNOLIA | SCARLET SAGE |
| AZALEA | ENGLISH IVY | MAPLE | SNAPDRAGON |
| BABYSBREATH | EUONYMUS | MARIGOLD | SPIREA |
| BARBERRY | EVENING PRIMROSE | MOCK ORANGE | SPRUCE |
| BELLFLOWER | FAILLARDIA | MORNING GLORY | STONE CROP |
| BIRCH | FERNLEAF YARROW | MOTHER OF THYME | STRAWFLOWER |
| BLEEDING HEART | FEVERFEW | MOUNTAIN LAUREL | SUNFLOWER |
| BLOOD LEAF | FIG | MOURNING BRIDE | SUNFLOWER |
| BOXWOOD | FORGET ME NOT | MOURNING PINKS | SWEET PEA |
| BUGLOSS | FORSYTHIA | NASTURTIUM | SYCAMORE |
| CAMELLIA | FOUR O'CLOCK | OAK | THREE PEONY |
| CANDLE LARKSPUR | FOXGLOVE | ORPINE | TULIP TREE |
| CANDYTUFTS | GERANIUM | PACHISTIMA | VARIEGATED PRIVET |
| CHESTNUT | GLADIOLUS | PACHYSANDRA | VERBENA |
| CHRYSANTHEMUM | GOLD DUST | PEONIES | VIOLET |
| CINQUEFOIL | GOLDEN MARGUERITE | PETUNIA | VIBURNUM |
| COLEUS | GOLDEN TUFT | PINE | VIRGINIA SPIDERWORT |
| COLUMBINE | GUM | PINK HEATH | WALNUT |
| CORAL BELLS | HAWTHORN | PITTOSPORUM | WEIGELA |
| COREOPSIS | HOLLY | PODOCARPUS | WILLOW |
| COSMOS | HONEYSUCKLE | POKER PLANT | WILSON IVY |
| COTONEASTER | HYDRANGEA | POPLAR | WORMWOOD |
| COTTONWOOD | IRIS | PRIVET | YEW |
| CRABAPPLE | JAPANESE ANDROMEDA | PURPLE CONEFLOWER | ZINNIA |
| CUPHEA | JUNIPER | | |

SHOULD NOT BE USED ON THE FOLLOWING PLANTS: Bugle Weed, Button Pink, Carnation, Guar, Germander, Mesembryanthemum, Pansy, Phlox, Sweet William and Tradescantia.

LIST OF WEEDS CONTROLLED

- | | | |
|------------------------------|---------------------------|--------------------------|
| ANNUAL BLUEGRASS | GREEN FOXTAIL | PURSLANE |
| BROWN TOP PANICUM | JOHNSON GRASS (From Seed) | REEDROOT PIGWEED |
| CARPETWEED | LAMBSQUARTER | SESBANIA (S. Microcarpa) |
| CHICKWEED (Common) | LOVEGRASS | TEXAS MILLET |
| CRABGRASS (Hairy and Smooth) | NODDING SPURGE | WITCHGRASS |
| FLORIDA PUSSLEY | | YELLOW FOXTAIL |

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking. Do not store near feed foodstuffs, seeds, fertilizers or other pesticides. Store in a dry place. Do not graze on treated areas or feed plant refuse to livestock. Do not reuse container. Destroy when empty, either burying with other wastes away from water supplies, or by burning. Do not inhale smoke or fumes.

NOTE: Follow directions carefully as use, storage and handling of this material is beyond the control of the seller; the buyer assumes all risks. If directions given hereon are not strictly adhered to.

SOLD BY:

5535-102

EPA Reg. No. 5535-102