

Feb 13, 1975

5535

4M-10-74

GRO-WELL

DIAZINON*

25E

INSECT SPRAY

SINCE 1855

**FOR CONTROL OF CERTAIN INSECTS ATTACKING
FLOWERS, SHRUBS, SHADE TREES AND
CERTAIN FRUITS AND VEGETABLES
MAKES UP TO 48 GALLONS OF SPRAY**



ACTIVE INGREDIENTS:	
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....	25.00%
Aromatic Petroleum Derivative Solvent.....	57.00%
INERT INGREDIENTS:	18.00%
TOTAL	100.00%

*DIAZINON® Trademark of CIBA GEIGY CORP.

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS
NET CONTENTS: 16 FL. OZ. (1 PINT)**

EPA EST. NO 904-NJ-1

GRO-WELL® DIAZINON 25E INSECT SPRAY will control the insects listed below which attack **ROSES, FLOWERS, SHRUBS, TREES, LAWNS, VEGETABLES** and **FRUITS**. A sprayer which attaches to a hose, such as the **GRO-WELL SHRUB & TREE SPRAYER** or **GRO-WELL LAWN SPRAYER** (for Lawn and Vegetable areas), may be used. Read entire label before using. Use strictly in accordance with label cautions, warnings and directions. Do not store diluted spray. Make a new dilution for each use, following ratios listed in **DILUTION** column, mixing the fluid ounces with the gallons of water as noted. Apply when insects appear and repeat as noted in directions, or as necessary. For lawns and vegetable gardens measure area to be treated and apply to the square foot area as indicated below.

FOR SPRAYING	TO CONTROL	DILUTION		AREA COVERAGE
		CHEMICAL (FL. OZ.)	WATER (GALS.)	
LAWN SPRAY ¹	Cinch Bugs, Sod Webworms, Ants ⁹ , Ticks, Earwigs	4	12	500 sq. ft. (25' x 20')
	White Grubs ² (Japanese Beetle Larvae and European Chafer), Billbugs ³	4	5-15	500 sq. ft. (25' x 20')
	Hyperodes Weevils ⁴ (New York State)	3	5-15	500 sq. ft.
FLOWER, SHRUB & SHADE TREE SPRAY ⁵ Roses, Azaleas, Chrysanthemum, Carnations, Gladioli, Oaks, Maples, Locust, Holly, Boxwood, Pine, Spruce, Willow	Aphids, Mites, Leafhoppers, Thrips, Leafminers, Scale Crawlers	1	3	
FRUIT SPRAY ⁶ Apples, Peaches, Plums, Cherries, Grapes, Strawberries	Codling Moths ¹⁰ , Mites, Aphids, Leafrollers, Scale Crawlers	4	12	
VEGETABLE SPRAY ⁷ Tomatoes, Carrots, Potatoes, Beans, Onions, Cucumbers, Cantaloupe, Cabbage, Peppers	Bean Beetles, Leafminers, Aphids, Cucumber Beetles, Imported Cabbageworms	1	3	
PRIOR TO PLANTING ⁸ Cabbage, Carrots, Peas, Lettuce, Radishes, Snapbeans, Tomatoes	Cutworms (Surface and Subterranean), Wireworms, Garden Symphylans, Root Maggots	7½	12	500 sq. ft. (25' x 20')
OUTDOOR AREA ⁹ Nuisance Pests	Flies, Ants, Ticks, Boxelder Bugs, Cockroaches, Clover Mites, Spiders, Earwigs, Chiggers	8	1½	

NOTES: 1. Apply a few hours after a heavy rain or thorough watering for control of all listed lawn insects. 2. Apply between late July and early October. Water thoroughly after application. 3. Apply when insects or damage first appear. 4. Apply first treatment in mid-April, repeat in mid-May. 5. Penetrate dense foliage by spraying the upper and lower surfaces of the foliage thoroughly. 6. Start spraying after blooming and repeat at 10-14 day intervals if needed. Do not spray within 21 days of harvest. 7. Three (3) gallons of diluted spray will cover 1,000 sq. ft. (50' x 20'). Spray at 7-10 day intervals when insects appear. Do not spray within 14 days of harvest, however, tomatoes may be sprayed 1 day before harvest. Wash home garden vegetables before using. 8. Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix spray into the top 4-8 inches of soil. 9. Thoroughly spray window frames, doors, sills, screens, porches, patio walls, foundations, walks, garbage cans and into cracks and other places where insects may hide, especially ants, paying particular attention to their hills and trails. Keep small children away from treated areas. Do not treat plants with this dosage rate. 10. Apply ¹⁰ metal fall and every 10 to 14 days thereafter until control is achieved.

WARNING: KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. MAY BE FATAL IF SWALLOWED. FOR OUTDOOR USE ONLY. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on humans or household pets. Keep children and pets off treated areas until spray has completely dried. Do not use, pour, spill or store near heat or open flame.

ENVIRONMENTAL CAUTION: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use on certain Ferns, Poinsettias, Hibiscus, Papaya, Pilea and Gardenia, since injury to these plants may occur.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given herewith are not strictly adhered to.

SOLD BY: J. & L. ADIKES, INC., JAMAICA, N. Y. 11423 EPA Reg. No. 5535 -103

Feb 13, 1975

5535

4M-10-74

GRO-WELL

DIAZINON*

25E

INSECT SPRAY

SINCE 1855

**FOR CONTROL OF CERTAIN INSECTS ATTACKING
FLOWERS, SHRUBS, SHADE TREES AND
CERTAIN FRUITS AND VEGETABLES
MAKES UP TO 48 GALLONS OF SPRAY**



ACTIVE INGREDIENTS:	
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....	25.00%
Aromatic Petroleum Derivative Solvent.....	57.00%
INERT INGREDIENTS:	18.00%
TOTAL	100.00%

*DIAZINON® Trademark of CIBA GEIGY CORP

**WARNING: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS
NET CONTENTS: 16 FL. OZ. (1 PINT)**

GRO-WELL® DIAZINON 25E INSECT SPRAY will control the insects listed below which attack **ROSES, FLOWERS, SHRUBS, TREES, LAWNS, VEGETABLES** and **FRUITS**. A sprayer which attaches to a hose, such as the **GRO-WELL SHRUB & TREE SPRAYER** or **GRO-WELL LAWN SPRAYER** (for Lawn and Vegetable areas), may be used. Read entire label before using. Use strictly in accordance with label cautions, warnings and directions. Do not store diluted spray. Make a new dilution for each use, following ratios listed in **DILUTION** column, mixing the fluid ounces with the gallons of water as noted. Apply when insects appear and repeat as noted in directions, or as necessary. For lawns and vegetable gardens measure area to be treated and apply to the square foot area as indicated below.

FOR SPRAYING	TO CONTROL	DILUTION		AREA COVERAGE
		CHEMICAL (FL. OZ.)	WATER (GALS.)	
LAWN SPRAY ¹	Cinch Bugs, Sod Webworms, Ants ² , Ticks, Earwigs	4	12	500 sq. ft. (25' x 20')
	White Grubs ² (Japanese Beetle Larvae and European Chafer), Billbugs ³	4	5-15	500 sq. ft. (25' x 20')
	Hyperodetes Weevils ⁴ (New York State)	3	5-15	500 sq. ft.
FLOWER, SHRUB & SHADE TREE SPRAY ⁵ Roses, Azaleas, Chrysanthemum, Carnations, Gladiolus, Oaks, Maples, Locust, Holly, Boxwood, Pine, Spruce, Willow	Aphids, Mites, Leafhoppers, Thrips, Leafminers, Scale Crawlers	1	3	
FRUIT SPRAY ⁶ Apples, Peaches, Plums, Cherries, Grapes, Strawberries	Codling Moths ¹⁰ , Mites, Aphids, Leafrollers, Scale Crawlers	4	12	
VEGETABLE SPRAY ⁷ Tomatoes, Carrots, Potatoes, Beans, Onions, Cucumbers, Cantaloupe, Cabbage, Peppers	Bean Beetles, Leafminers, Aphids, Cucumber Beetles, Imported Cabbageworms	1	3	
PRIOR TO PLANTING ⁸ Cabbage, Carrots, Peas, Lettuce, Radishes, Snapbeans, Tomatoes	Cutworms (Surface and Subterranean), Wireworms, Garden Symphylans, Root Maggots	7½	12	500 sq. ft. (25' x 20')
OUTDOOR AREA ⁹ Nuisance Pests	Flies, Ants, Ticks, Boxelder Bugs, Cockroaches, Clover Mites, Spiders, Earwigs, Chiggers	8	1½	

EPA EST. NO 904-NJ-1

NOTES: 1. Apply a few hours after a heavy rain or thorough watering for control of all listed lawn insects. 2. Apply between late July and early October. Water thoroughly after application. 3. Apply when insects or damage first appear. 4. Apply first treatment in mid-April, repeat in mid-May. 5. Penetrate dense foliage by spraying the upper and lower surfaces of the foliage thoroughly. 6. Start spraying after blooming and repeat at 10-14 day intervals if needed. Do not spray within 21 days of harvest. 7. Three (3) gallons of diluted spray will cover 1,000 sq. ft. (50' x 20'). Spray at 7-10 day intervals when insects appear. Do not spray within 14 days of harvest, however, tomatoes may be sprayed 1 day before harvest. Wash home garden vegetables before using. 8. Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix spray into the top 4-8 inches of soil. 9. Thoroughly spray window frames, doors, sills, screens, porches, patio walls, foundations, walks, garbage cans and into cracks and other places where insects may hide, especially ants, paying particular attention to their hills and trails. Keep small children away from treated areas. Do not treat plants with this dosage rate. 10. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

WARNING: KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. MAY BE FATAL IF SWALLOWED. FOR OUTDOOR USE ONLY. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on humans or household pets. Keep children and pets off treated areas until spray has completely dried. Do not use, pour, spill or store near heat or open flame.

ENVIRONMENTAL CAUTION: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Pilea and Gardenia, since injury to these plants may occur. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller. The buyer assumes all risks if directions given herewith are not strictly adhered to.

SOLD BY: J. & L. ADIKES, INC., JAMAICA, N. Y. 11423 EPA Reg. No. 5535 103

GRO-WELL® DIAZINON 25E INSECT SPRAY will control the insects listed below which attack **ROSES, FLOWERS, SHRUBS, TREES, LAWNS, VEGETABLES** and **FRUITS**. A sprayer which attaches to a hose, such as the **GRO-WELL SHRUB & TREE SPRAYER** or **GRO-WELL LAWN SPRAYER** (for Lawn and Vegetable areas), may be used. Read entire label before using. Use strictly in accordance with label cautions, warnings and directions. Do not store diluted spray. Make a new dilution for each use, following ratios listed in **DILUTION** column, mixing the fluid ounces with the gallons of water as noted. Apply when insects appear and repeat as noted in directions, or as necessary. For lawns and vegetable gardens measure area to be treated and apply to the square foot area as indicated below.

FOR SPRAYING	TO CONTROL	DILUTION		AREA COVERAGE
		CHEMICAL (FL. OZ.)	WATER (GALS.)	
LAWN SPRAY ¹	Cinch Bugs, Sod Webworms, Ants ⁹ , Ticks, Earwigs	4	12	500 sq. ft. (25' x 20')
	White Grubs ⁷ (Japanese Beetle Larvae and European Chafer), Billbugs ³	4	5-15	500 sq. ft. (25' x 20')
	Hyperodades Weevil ⁴ (New York State)	3	5-15	500 sq. ft.
FLOWER, SHRUB & SHADE TREE SPRAY ⁵ Roses, Azaleas, Chrysanthemum, Carnations, Gladioli, Oaks, Maples, Locust, Holly, Boxwood, Pine, Spruce, Willow	Aphids, Mites, Leafhoppers, Thrip, Leafminers, Scale Crawlers	1	3	
FRUIT SPRAY ⁶ Apples, Peaches, Plums, Cherries, Grapes, Strawberries	Codling Moths ¹⁰ , Mites, Aphids, Leafrollers, Scale Crawlers	4	12	
VEGETABLE SPRAY ⁷ Tomatoes, Carrots, Potatoes, Beans, Onions, Cucumbers, Cantaloupe, Cabbage, Peppers	Beet Beetles, Leafminers, Aphids, Cucumber Beetles, Imported Cabbageworms	1	3	
PRIOR TO PLANTING ⁸ Cabbage, Carrots, Peas, Lettuce, Radishes, Snapbeans, Tomatoes	Cutworms (Surface and Subterranean), Wireworms, Garden Symphylans, Root Maggots	7½	12	500 sq. ft. (25' x 20')
OUTDOOR AREA ⁹ Nuisance Pests	Fires, Ants, Ticks, Boxelder Bugs, Cockroaches, Clover Mites, Spiders, Earwigs, Chiggers	8	1½	

NOTES: 1. Apply a few hours after a heavy rain or thorough watering for control of all listed lawn insects. 2. Apply between late July and early October. Water thoroughly after application. 3. Apply when insects or damage first appear. 4. Apply first treatment in mid April, repeat in mid May. 5. Penetrate dense foliage by spraying the upper and lower surfaces of the foliage thoroughly. 6. Start spraying after blooming and repeat at 10-14 day intervals if needed. Do not spray within 21 days of harvest. 7. Three (3) gallons of diluted spray will cover 1,000 sq. ft. (50' x 20'). Spray at 7-10 day intervals when insects appear. Do not spray within 14 days of harvest, however, tomatoes may be sprayed 1 day before harvest. Wash home garden vegetables before using. 8. Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix spray into the top 4-8 inches of soil. 9. Thoroughly spray window frames, doors, sills, screens, porches, patio walls, foundations, walks, garbage cans and into cracks and other places where insects may hide, especially ants, paying particular attention to their hills and trails. Keep small children away from treated areas. Do not treat plants with this dosage rate. 10. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

WARNING: KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. MAY BE FATAL IF SWALLOWED. FOR OUTDOOR USE ONLY. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on humans or household pets. Keep children and pets off treated areas until spray has completely dried. Do not use, pour, spill or store near heat or open flame.

ENVIRONMENTAL CAUTION: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Pilea and Gardenia, since injury to these plants may occur.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

EPA EST. NO 904-NJ-1



OTHER

GRO-WELL

PRODUCTS

- MULTI-PURPOSE DUST
- CHLORDANE
- LAWN FUNGICIDE
- LAWN WEED KILLER
- CRABGRASS KILLER
- INDOOR-OUTDOOR AEROSOL SPRAY
- ROSE AEROSOL SPRAY
- DORMANT SPRAY
- FRUIT TREE SPRAY
- ROSE DUST
- TOMATO & VEGETABLE DUST
- MALATHION
- TREE WOUND PAINT
- HORNET BOMB
- CHICKWEED KILLER
- GRO-WELL LAWN SEED
- LAWN FOOD
- ACID PLANT FOOD
- GARDEN PLANT FOOD
- ROSE FOOD

SOLD BY: J. & L. ADIKES, INC., JAMAICA, N. Y. 11423 EPA Reg. No. 5535 - 103