


50 LBS. NET

5% SEVIN[®] DUST



ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate)

INERT INGREDIENTS:

5%
95%
TOTAL 100%

CAUTION

Keep Out Of Reach Of Children

READ THE LABEL

Caution: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid breathing of dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothes daily. Avoid storage near feed and food products.

Note for Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidote.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE OF WARRANTY

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

GENERAL DIRECTIONS

Apply Sevin Dust when first signs of insect infestation are noted, preferably before damage from feeding occurs. For best results, apply thoroughly to upper and lower leaf surfaces. If, needed, repeat application every 7 to 10 days.

Sevin Dust can be used as directed and applied to beans the day of harvest without exceeding the residue tolerance.

Sevin Dust is compatible with commonly used miticides, fungicides, and insecticides. Do not use with lime or other alkaline material. Sevin dust will not control insects if used with lime or other highly alkaline materials.

DIRECTIONS FOR USE

Beans: To control Mexican Bean Beetle, Bean Leaf Beetle, Corn Earworm, Leafhoppers, Spotted Cucumber Beetle. Apply 25 to 30 lbs. per acre.

Potatoes: To control Colorado Potato Beetle, Flea Beetle, Leafhopper. Apply 25 to 30 lbs. per acre.

Sweet Corn and Field Corn: To control Corn Earworm and Fall Army Worm. Apply at rate of 30 to 40 lbs. per acre. Begin dusting when first silks appear and repeat at 2 to 3 day intervals. Do not apply within 7 days of harvest for forage. Application of this product to the tassel region of sweet corn during the pollen shed period will seriously reduce bee populations.

Cucumbers, Summer Squash and Cantaloupes: To control Spotted Cucumber Beetle, Squash Bug, Flea Beetle. Apply at rate of 20 lbs. per acre. To control Melon Worm and Pickleworm. Apply at rate of 15 to 20 lbs. per acre. Thorough coverage is essential. May be applied up to harvest. Make only one application on cucumbers because of possible foliar injury.

Tobacco: Hornworms, Budworms, Japanese Beetle, Flea Beetles. Apply 15 to 25 lbs. per acre. Good coverage is essential, especially for Flea Beetle control. Pay particular attention to underside of leaves over entire plant.

— Manufactured By —

Coastal Chemical Corp.

Greenville, N. C.

EPA Reg. No. 5549-10

*Trade Mark of Union Carbide Corporation for the active ingredient — 1-naphthyl N-methylcarbamate.