




50 LBS. NET

4% Copper

5% Sevin - 75% Sulphur

Peanut Dust



**ACTIVE INGREDIENTS:**

- Cuprous Oxide
- (3.55% Metallic Copper)
- Carbaryl (1-Naphthyl N-Methylcarbamate)
- Sulphur

**INERT INGREDIENTS:**

	4%
	5%
	75%
	16%
<b>TOTAL</b>	<b>100%</b>

**DIRECTIONS FOR USE**

**LEAFSPOT CONTROL:** Apply 15 to 20 lbs. per acre. Start dusting during the last week of June and continue applications at two week intervals for a total of three to four applications. Any application washed off by rain within 24 hours should be repeated as soon as the weather is settled.

**INSECT CONTROL:** Spotted Cucumber Beetle, Southern Army Worm, Corn Earworm, Fall Army Worm, Thrips, apply 20 lbs. per acc. Repeat any application washed off by rain within 24 hours.

**CAUTION**

**Keep Out Of Reach Of Children**

Avoid contact with skin and eyes. Avoid inhalation of dust. Keep clothing free from dust. Harmful if swallowed, or inhaled.

Avoid storage near feed and foodstuffs. Wash thoroughly after handling. For protection of honey bees avoid use, if possible during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

\*Trade Mark of UCC for Carbaryl (1-Naphthyl N-Methylcarbamate)  
U. S. Patent No. 2,903,478

**NOTICE**

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured By

**COASTAL CHEMICAL CORPORATION**

GREENVILLE, N. C.