



# RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



## Parathion <sup>MAY 31 1978</sup> Dust

### ACTIVE INGREDIENT:

Parathion (O,O-Diethyl O-p-nitrophenyl phosphorothioate)

### INERT INGREDIENTS:

Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, for the pesticide registered under EPA Reg. No. 5549-304

5-30-78

1.0%

99.0%

Total 100.0%

### DIRECTIONS FOR USE

**TOBACCO PLANT BED:** For control of Flea Beetle, Midge & Crane Fly larva, Vegetable Weevil, Aphid, Green June Beetle larva, Suckfly, Split worm. Use 1/2 to 1 1/2 lbs. to 100 square yards. Make 1 to 4 applications as required. Newly set tobacco plants: Use 6 to 10 lbs. per acre. Treat entire plant. Larger tobacco plants in field use 10 to 25 lbs. per acre. Apply dust thoroughly to all parts of plants. Repeat if necessary. Avoid heavy use just prior to priming or cutting. Make no application within 5 days of priming or 15 days of cutting. Avoid plant juices coming in contact with skin of those engaged in cutting or priming tobacco.

**COTTON:** Red Spiders—apply 25 lbs. per acre. Make 2 to 3 applications at 5 to 7 day intervals when leaves first begin turning yellow or reddish brown. Aphids—apply 25 lbs. per acre when honeydew first appears and leaves begin to curl downward. Do not apply to cotton within 7 days of hand picking. Workers entering fields within 24 hours of treatment should wear protective clothing. Do not apply within 15 days of harvest if cotton trash is to be fed to dairy animals or animals being finished for slaughter.

**CABBAGE:** Use 40 to 50 lbs. per acre for control of diamond-back moth, Imported Cabbage Worm, Army Worm, Aphids & Thrips. Do not apply within 21 days of harvest.

**BEANS:** Use 40 lbs. per acre for control of Mexican Bean Beetles, Leaf-Hoppers, Aphids, Red Spiders, Army Worms, Leaf Rollers and Leaf Miners. Apply at 7 to 10 day intervals as required. Do not apply within 21 days of harvest.

**WARNING:** May be fatal if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear mask or respirator of a type by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and food-stuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of dust from crop areas treated. Birds and other wildlife in treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

### NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

—MANUFACTURED BY—

# Coastal Chemical Corporation

25 LBS. NET

EPA Reg. No. 5549-24

GREENVILLE, N. C.

EPA Est. 5549-NC-1

BEST DOCUMENT AVAILABLE