

[142]

MALATHION

88-101-1


Active ingredient:
Malathion
Inert ingredients

100
100

Use directions and disposal of this product as directed.

WARNING

Keep out of Reach of Children



CAUTION: Avoid prolonged breathing of dust. Avoid repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and food stuffs. Do not apply under conditions involving possible drift to food, forage or other plantings that may be damaged or the reproductive period of plants until for sale, use or consumption.

This product is toxic to fish. Keep out of water bodies or water which may be used for drinking water. Do not contaminate water by cleaning of equipment or disposal. Do not use in or near fish ponds or other aquatic areas.

Use entire label as a guide. Do not mix with oil, grease, kerosene and kerosene. Do not mix with ammonia. State of Federal Registration. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

DO NOT MIX

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

Use only as directed. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions. Do not use in areas where it is prohibited by local weather conditions.

[292]

Campana's Note
Poor Copy

Ornamental: Aphids, Red Spider Mites, Whiteflies, Japanese Beetle, Adult, Four-lined Leaf Bug, Bagworm, Carnation Leaf Miner, Rose Leafhopper, Potato Leafhopper. Repeat as necessary. Do not use on Boston Maiden hair and Pteris ferns, and some species of grasses.

Peas: Pea Aphid. Apply as needed. Do not apply within 7 days of harvest if pea forage is to be fed to livestock.

Potatoes: Aphids, Leafhoppers. Use or not application as needed. Use up to harvest.

Strawberries: Aphids, Spider Mites, Lygus Bugs, Flea Beetles, Strawberry Leafhopper, Strawberry Leafrollers, Whiteflies, Strawberry Root Weevil, Field Crickets. Repeat as necessary. Strawberry Root Weevil. Apply to soil surface before planting and work into top 2 to 3 inches of soil. Tobacco: Aphids. Make full coverage application. Repeat as needed. Do not apply within 7 days of harvest.

Tomatoes: Aphids, Spider Mites, Lygus Bugs, Flea Beetles, Strawberry Root Weevil. Use 1 lb. per acre. Do not apply within 7 days of harvest.

Billipedes, principally lawns in and around greenhouses, around houses, in yards and gardens where beetles and other insects are known to be present. Use 1 lb. per 1,000 sq. ft. of soil surface or where insects congregate. Repeat at 7 to 10 day intervals as needed.

Fleas: In dog and cat kennels, in pens, yards, lawns, and under houses. Use 1 to 2 lbs. per 1,000 sq. ft. Use a hand lister. Remove manure or debris before dusting.

Poultry Houses: Use as needed. Use as needed. Mites, fleas, lice. Apply thoroughly to roosts, feed troughs, and areas where birds roost. Litter at 1 lb. per 1,000 sq. ft. of floor space. Repeat as necessary. Avoid contamination of feed and water.

Poultry (Direct application to birds): Lice, fleas, mites, and ticks. Northern Fowl Mites, Poultry Lice, Chicken Red Mite. Use as supplement to premise treatment for Chicken Red Mite. Apply dust to birds thoroughly. Apply dust by means of shaker can or spray hand dusting. Repeat in 4 to 6 weeks or when necessary. Also treat all birds in contact.

NEED FOR A...

River assays and field use, should be made in accordance with the following instructions in strict accordance with directions for use.

MANUFACTURE...

MANUFACTURE...

MANUFACTURE...

MANUFACTURE...

MANUFACTURE...