




50 LBS. NET

# 20% TOXAPHENE DUST



## ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene—containing 67-69% chlorine) 20.00%

INERT INGREDIENTS: 80.00%

TOTAL 100.00%

## DIRECTIONS

**COTTON:** Boll Weevil, Boll Worm, Cotton Fleahopper, Tarnished Plant Bug, Rapid Plant Bug, Cotton Leafworm, Thrips, Fall Armyworm, Grasshopper - Apply 10 to 20 lbs. per acre, depending on degree of infestation and stage of plant growth. Begin application at 5 day intervals when one or more adult weevils are found per 100 terminals one week before squaring (3-10 leaf stage), or the square infestation first rises to 10% if pre-square application were not needed. Continue applications as long as the plants are squaring.

**TOMATOES:** Fruit Worm, Pinworm, Hornworm, Blister Beetle - Apply 10 to 20 lbs. per acre. Do not apply within 7 days of harvest.

**BEANS:** Mexican Bean Beetle, Thrips - Apply 10 to 20 lbs. per acre. Do not apply to Green or String Beans within 7 days of harvest. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**SOYBEANS:** Fall Armyworm, Green Cloverworm, Velvet-Bean Caterpillar, Grasshopper, Corn Earworm, Bean Leaf Beetle - Use 10 to 20 lbs. per acre. Do not treat soybeans within 21 days of harvest. Do not feed treated plant or ensilage made from treated plants to poultry, dairy animals and animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

## CAUTION

### Keep Out Of Reach Of Children

Harmful if swallowed. Avoid breathing of dust mist. Avoid contact with eyes, skin, and clothing. Wash thoroughly after using. Avoid contamination of feeds or foodstuffs. Keep away from children and domestic animals. Do not pasture dairy animals being fattened for slaughter in treated fields. Do not use on cucurbites.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**ANTIDOTE:** INTERNAL - Give sodium chloride 15 g. (one tablespoonful of table salt) in a glass of warm water and repeat until vomit is clear. Call a doctor immediately. EXTERNAL - Wash with soap and water.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

## NOTICE

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

— Manufactured By —

EPA Reg. No. 5549-61

**Coastal Chemical Corp.**

GREENVILLE, N. C.