

CAUTION

Do not use treated crops for feed or pasturing of poultry, dairy animals, or animals being finished for slaughter.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Birds and other wildlife in treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of waste.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not reuse the empty container. Destroy by perforating or crushing. Bury or discard in a safe place.

WARNING

May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using. Avoid contamination of food, food containers and feedstuffs. Do not use this product on dairy animals, in dairy barns or on feed forage to be fed to dairy animals or animals being finished for slaughter.

Toxaphene Emulsion

Contains 6 lbs of Toxaphene per gallon

ACTIVE INGREDIENTS	BY WEIGHT
* Toxaphene	60.00%
Xylene	35.00%
INERT INGREDIENTS	5.00%
Total	100.00%

* Technical Chlorinated Camphene (Chlorine Content 67% to 69%).

WARNING

Keep Out Of Reach Of Children

See Side Panel for additional Warnings

— **Manufactured By** —

Coastal Chemical Corporation

GREENVILLE, N. C.

Net Contents — 1 Gallon

EPA Reg. No. 5549-62

DIRECTION

COMMERCIAL ORNAMENTAL: Use 1 1/2 to 2 quarts per 1000 sq. ft. Treated areas should be kept free of children and pets off treated areas. If the insecticide has been washed off, grass has completely dried.

BEANS: Mexican Bean, Earworm: Use 1 1/2 to 2 quarts per acre. Use water for good coverage. Do not apply to beans after the pods start to form.


Do not feed treated beans to animals being finished for slaughter.

SOYBEANS: Armyworm, Corn Earworm, Grasshopper: Use 1 1/2 to 2 quarts per acre. Do not apply to soybeans after the pods start to form.

Do not feed treated plants to poultry, being finished for slaughter. Do not use mill trash to livestock or poultry.

NOTICE

Buyer assumes all risks of use of this material not in strict accordance with directions given herewith.



ION

for feed or pasturing of animals being finished

fish and wildlife. Keep out of water. Do not apply where runoff may reach other wildlife in treated areas.

to bees exposed to direct spray. Protective information from your Cooperative Agricultural Agent.

Under conditions favoring drift, do not contaminate water by spraying or disposal of waste.

DO NOT SPILL OR STORE NEAR FEED.

Destroy by pouring contents into water or discard in a safe place.

WARNING

Do not breathe spray, get on skin, or on clothing. Avoid contamination of feedstuffs. Do not use in dairy barns or on dairy animals or animals being finished.

Toxaphene Emulsion

Contains 6 lbs of Toxaphene per gallon

ACTIVE INGREDIENTS	BY WEIGHT
• Toxaphene	60.00%
Xylene	35.00%
INERT INGREDIENTS	5.00%
Total	100.00%

• Technical Chlorinated Camphene (Chlorine Content 67% to 69%).

WARNING
Keep Out Of Reach Of Children

See Side Panel for additional Warnings

— **Manufactured By** —

Coastal Chemical Corporation

GREENVILLE, N. C.

Net Contents — 1 Gallon

EPA Reg. No. 5549-62

DIRECTIONS FOR USE

COMMERCIAL ORNAMENTAL TURF. Sod Webworm, Armyworms, Fall Armyworms: Use 6 to 8 oz. per 1000 sq. ft. Treat in late afternoon. Keep children and pets off treated areas until the insecticide has been washed off by sprinkling and the grass has completely dried.

BEANS: Mexican Bean Beetle, Leafhoppers, Corn Earworm: Use 1 1/2 to 2 quarts per acre in sufficient water for good coverage. Do not treat green or snap beans after the pods start to form.

Do not feed treated bean vines to dairy animals or animals being finished for slaughter.

SOYBEANS: Armyworm, Velvet Bean Caterpillar, Corn Earworm, Grasshoppers: Mix 1 1/2 to 2 2/3 quarts with sufficient water to give good coverage per acre. Do not apply within 21 days of harvest.

Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

