

SEVIN* MALATHION DUST

ACTIVE INGREDIENTS:

CARBARYL (1-NAPHTHYL N-METHYLCARBAMATE)	3%
MALATHION (O, O-DIMETHYL DITHIOPHOSPHATE OF DIETHYL MERCAPTOSUCCINATE)	3%
INERT INGREDIENT	94%
TOTAL	100%

CAUTION

Keep Out Of Reach Of Children

READ THE LABEL

CAUTION: Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Note to Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, feeding troughs and watering receptacles. This product is toxic to fish. Keep out of any body of water. Fish may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE OF WARRANTY

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured By

Coastal Chemical Corporation

Greenville, N. C.

EPA Reg. No. 5549-66

Net Weight 25 lbs.

*Trade mark of Union Carbide Corporation for the active ingredient — 1-naphthyl N-methylcarbamate.

DIRECTIONS FOR USE

Begin Application when insects or damage is first observed. Apply with a good dust gun or blower so as to give good coverage to both upper and lower surface of the leaves. Repeat application on a 5 to 7 day schedule or as necessary for good insect control.

BEANS: To control Bean Leaf Beetle, Corn Earworm, Leafhopper, Mexican Bean Beetle, Red Spider Mites, Cucumber Beetles, Japanese Beetle, Lygus Bugs, Flea Beetle and Stink Bug: Use 34 pounds per acre. To control Aphids: Use 50 pounds per acre.

CABBAGE, COLLARDS: To control Aphids, Cabbage Looper: Use 50 pounds per acre. To control Harlequin Bug, Imported Cabbageworm, Corn Earworm and Armyworm: Use 34 pounds per acre.

ENGLISH PEAS: To control Aphids, Bean Leaf Beetle and Pea Weevil: Use 34 pounds per acre.

OKRA: To control Aphids, Corn Earworm, and Stink Bug: Use 34 pounds per acre. Do not apply after the pods begin to form.

IRISH POTATOES: To control Colorado Potato Beetle, Flea Beetle, Aphids, Leafhopper, False Chinch Bug: Use 42 pounds per acre.

TOMATOES, EGGPLANTS, PEPPERS: To control Flea Beetle, Tomato Fruitworm, Tomato Hornworm, European Corn Borer, Stink Bug, and Fall Armyworm: Use 34 pounds per acre. To control Red Spider Mites, Aphids, Tomato Russet Mite: Use 50 pounds per acre. To control Leafhopper: Use 67 pounds per acre.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKIN, MELONS: To control Cucumber Beetle, Flea Beetle, Leafhopper: Use 34 pounds per acre. To control Pickworm, Melonworm, Aphids and Spider Mites: Use 50 pounds per acre. Make only one application on cucumbers due to possible foliar injury. Do not apply to cucurbits unless plants are dry.

Interval Before Harvest: Can be used up to day of harvest on Irish Potatoes; one day on Beans, Tomatoes, Cucumbers, Squash, Cantaloupe, Melons; Three days on English Peas, Eggplants, Peppers, Pumpkins; seven days on Cabbage; fourteen days on Collards.

NOTICE: Make no application within seven days if vines or leaves are to be used for animal feeds.

REGISTERED TRADE MARK
UNION CARBIDE CORPORATION
GREENVILLE, N. C.