

[5549-69]



# THURI-90 GARDEN DUST

## RECOMMENDATIONS

Thorough coverage essential when using THURI - 90 GARDEN DUST

CROP	PEST	LBS. PER ACRE
Artichoke	Artichoke plume moth	20 60
Broccoli	Cabbage looper	20 60
	Imported cabbage worm	16 50
Cabbage	Cabbage looper	20 60
	Imported cabbage worm	16 50
Cauliflower	Cabbage looper	20 60
	Imported cabbage worm	16 50
Celery	Cabbage looper	20 60
Collards	Cabbage looper	20 60
	Imported cabbage worm	16 50
Cucumbers	Cabbage looper	16 50
Grapes	Grape leaf folder	16 50
Kale	Cabbage looper	20 60
	Imported cabbage worm	16 50
Lettuce	Cabbage looper	20 60
Melons	Cabbage looper	16 50
Mustard Greens	Cabbage looper	20 60
	Imported cabbage worm	16 50
Potatoes	Cabbage looper	20 60
Tobacco	Tobacco hornworm	16 50
Tomatoes	Cabbager looper	20 60
	Tomato hornworm	16 50
Turnip Greens	Cabbage looper	20 60
	Imported cabbage worm	16 50

Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply by ground equipment using 16 50 pounds per acre to obtain thorough coverage of leaf surfaces.

**ACTIVE INGREDIENT:**  
 Bacillus thuringiensis Berliner, potency of 200 International Units (at least 375 thousand viable spores) per milligram — .04%  
**INERT INGREDIENTS:** \_\_\_\_\_ 99.96%  
**TOTAL** 100.00%

**CAUTION**  
 Keep Out Of Reach Of Children

Store in cool place.  
 For use only in southeastern United States (Alabama, Florida, Georgia, Kentucky, North Carolina, South Carolina, Tennessee, Virginia).  
 Container Disposal: Do not reuse empty container. Destroy or discard in safe place.

**NOTICE**  
 Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured By  
**Coastal Chemical Corp.**  
 Greenville, N. C.  
 Net Weight 25 lbs.

EPA Reg No. 5549-69

EPA Est. 5549-NC 1

**ACCEPTED**  
 MAY 17 1974  
 UNDER THE FEDERAL PESTICIDE  
 FUNGICIDE AND ROENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 5549-69

### Directions for Use

Apply recommended amounts when air is calm, using ground equipment, aircraft or hand duster. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Good underleaf coverage is essential since the insect must ingest Thuri - 90 Garden Dust. After eating Thuri - 90 Garden Dust, some insects stop feeding but may otherwise appear to be unaffected for several days.

Use the lowest recommendation only with hand duster and against light infestation.

Use the higher recommendations against very heavy infestations, or where it is difficult to obtain good coverage due to rank foliage. Thuri - 90 Garden Dust is exempt from tolerance requirements, and may be applied in recommended amounts up to the day of harvest.

