



DOUBLE DUTY DUST

Contains Sevin® and Dipel® Insecticides

Active Ingredients:

Bacillus thuringiensis, Berliner Potency of 240 International Units per mg... (at least 0.38 billion viable spores per g.)*	0.048%
Carbaryl (1-naphthyl N-methyl carbamate)	5.000%

Inert Ingredients:

Inert Ingredients	94.952%
-------------------	---------

*Sevin is a registered trademark of Union Carbide Corporation for carbaryl insecticide

*Dipel is a registered trademark of Abbott Laboratories.

*Equivalent to potency of 0.11 billion International Units per pound of this product

EPA Letter 5549-75

Total

100.000%

Precautionary Statements

Hazards to Humans

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not take internally. Avoid skin contact. Avoid getting in eyes. Wash hands before eating. Wash all contaminated clothing with soap and water before reuse. Avoid contamination of feedstuffs, feed troughs, and water receptacles. Wear gloves when mixing or applying this product.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. BEE CAUTION: May kill Honeybees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take other equally effective precautions.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Disposal

Do not reuse empty container. Wrap container and put in trash collection.

This product is a combination of Bacillus Thuringiensis, a biological insecticide, specifically for control of chewing caterpillars and Carbaryl, a broad spectrum insecticide, and will control the following insect pests on vegetables in the home garden: Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Leafhoppers, Flea Beetle, Japanese Beetle, Lygus Bug, Stink Bug, Lima Bean Pod Borer, Corn Earworm, Sap Beetle, Harlequin Bug, Armyworms, Imported Cabbageworm, Diamond-Back Moth Larvae, Cabbage Looper, Tarnished Plant Bug, Pickleworm, Melonworms, Squash Bug, European Corn Borer, Fall Armyworm, Spittlebug, Tomato Fruitworm, Tomato Hornworm, Colorado Potato Beetle.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- If Swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.
- If Inhaled: Remove victim to fresh air and apply respiration if indicated.
- If On Skin: Wash promptly with soap and water. Rinse thoroughly.
- If In Eyes: Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. See side panel for additional precautionary statements.

MANUFACTURED BY

Coastal Chemical Corporation

EPA Est. No. 5549-NC-1

EPA Reg. No. 5549-75

Net Contents-25 lbs.

DEC 10 1982

Sevin Insecticide Act
Dipel Insecticide Act
EPA Letter 5549-75

Directions for Use

Application: Apply dust at the rate of 2 oz. per 50 feet of row (40 lbs. per acre) at the first sign of insects using a good mechanical or hand-powered duster. Repeat as necessary for control. For best results, dust in early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. The number in () following each crop indicates days required between last application and harvest: (0) indicates it may be used up to and including day of harvest. This product may be used on the following vegetables: Beans (0), Broccoli (3), Brussels Sprouts (3), Cabbage (3), Cantaloupe (0), Carrots (0), Cauliflower (3), Collards (14), Corn (0), Cucumbers (0), Eggplants (0), Garden Beets (roots 3, tops 14), Kale (14), Kohlrabi (3), Lettuce (head 3, leaf 14), Mustard Greens (14), Okra (0), Peppers (0), Potatoes (0), Pumpkins (0), Radishes (3), Rutabaga (3), Squash (0), Spinach (14), Tomato (0), Turnips (root 3, tops 14).

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.