

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin or eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Do not apply directly on food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. Facilities and equipment in federally inspected meat and poultry plants must be washed with an effective cleaning compound followed by a potable water rinse after spraying. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, ponds, streams, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

**CONTAINER DISPOSAL**

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

**PESTICIDE DISPOSAL**

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to federal or approved state procedures under subtitle C of the Resource Conservation and Recovery Act.

**STORAGE**

Store in a cool dry area away from heat or open flame.

EPA REG. NO. 5576-52  
EPA EST. NO. 5576-Tx-1

DO NOT TRANSPORT OR STORE UNDER 30° F.

NET CONTENTS \_\_\_\_\_

FOR INDUSTRIAL, INSTITUTIONAL, OR PROFESSIONAL USE ONLY

**INSECTEND**

A Concentrated PYRENONE MicroEmulsion

-Clear- Effective Fast Acting Insecticide -Stable-

For Use in Homes, Institutions, Restaurants,

Food Processing Plants, Dairies, Livestock,

and Outdoors for Mosquito Control. Apply with

Conventional Mechanical or Compressed Air Equipment

**ACTIVE INGREDIENTS:**

Pyrethrins ----- 0.3%

\* Piperonyl Butoxide, Technical ----- 3.0%

Petroleum Distillate ----- 1.2%

INERT INGREDIENTS ----- 95.5%

100.0%

PYRENONE-Registered trademark of  
Fairfield American Corporation

\* Equivalent to 2.4% (butylcarbityl 6-propyl-  
piperonyl) ether and .60% related compounds.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

Harmful if swallowed. See side panel for  
additional precautionary statements.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**INDOOR APPLICATIONS**

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before use. vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SURFACE SPRAYING:** To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, drugstore beetles, rust red flour weevils, sawtoothed grain beetles, spider beetles, cigarette beetles, meal worms, grain mites and codellies: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored food, around the base of machinery, behind shelves and drawers. For silverfish spray bookcases, for ants, spray trails, nests, and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take best apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat as needed. To control fleas and brown dog ticks: Thoroughly spray infested areas, wet beds, resting quarters, nearby cracks and crevices along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air in sunlight for several hours. If possible, apply spray liberally to empty chests, closets, bureaus, and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least 3 feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog for rapid control of flies, fruit flies, gnats, mosquitoes, slipper flies, wasps, hornets, pompilids, grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000-40,000 cubic feet of space. Contact insects directly whenever possible.

**ON ANIMALS:** To protect cattle and horses from horn flies, house flies, mosquitoes, and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice: Apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 2-3 weeks if required.

**OUTDOOR APPLICATION**

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Low marshy places, stagnant pools or ponds and shorelines. Apply at a rate of 1/2 gallons per acre (1/2 ounces per 1000 square feet).

Using truck-mounted conventional misting equipment such as Sprayer Model 10000, 1000H traveling at 5 m.p.h. to treat a 300' x 100' area with 200' high vegetation, requires 1/2 gallons per minute.

This product is warranted to be of uniform quality within manufacturing tolerances. Since no control is exercised over its use, no warranty, expressed or implied, is made as to the effects of such use. The obligation hereunder shall be limited to refunding the purchase price of that portion of the material proven to be defective.

ACCEPTED

MAY 20 1985