


HUB STATES
KWIK-KILL FLY KRYSALS

K I L L S
Even Resistant

HOUSE FLIES



Active Ingredients:

2,2-Dichlorovinyl dimethyl phosphate5% *

Inert Ingredients

Total 100.0%

*Equivalent to .465% 2,2-Dichlorovinyl dimethyl phosphate
plus .035% insecticidally active related compounds.

Outdoor
Outdoor living areas * Backyard areas * Picnic grounds = Dog kennels = Ice Cream
Stands

Open Air Drive-Ins

Also effective for dairy barns * Milk sheds * Stables * Livestock barns = Poultry
Houses

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

SEE OTHER PANEL FOR ADDITIONAL PRECAUTIONS

HUB STATES CHEMICAL AND EQUIPMENT CORPORATION

2000 North Illinois Street

Indianapolis, Indiana 46202

DIRECTIONS FOR USE

Broadcast on the ground or floors in areas where flies gather or breed. Can be used in Barns, Stables and Loafing areas, feed rooms, poultry houses, kennels, around garbage disposals, manure piles and areas of certain commercial buildings where flies become a nuisance.

NOTICE: For best fly kill replace bait every 3 to 5 days.

DAIRY BARNS, STABLES, LIVESTOCK BARNS

Clean out manure and apply fly bait on clean floor areas or empty burlap sacks in areas frequented by flies. Treat areas outside barns where flies congregate. Use 1/4 lb. per 1,000 sq. ft. of area to be treated and repeat as necessary to reduce fly populations. Battery poultry establishments - scatter fly bait on droppings under cages, walkways and alleyways. For reducing flies in and around garbage collection and disposal areas and refuse areas where flies congregate. To make a spray add one-half pound of New Hub States Kwik-Kill Fly Krystals

to one gallon of water. To make bait stations place paper towels or other absorbent material such as burlap in squares, in pie plates or on boards and saturate with new Hub States Kwik-Kill Fly Krystals. After flies have been brought under control treatment should be repeated as needed. Do not use as a spray in dairy barns.

FACE FLY ON DAIRY AND BEEF CATTLE

To make a bait dilute 7/8 cup of warm water plus 1/2 cup warm sugar or corn syrup. Since this mixture loses the desired strength, mix only the amount needed for daily application. For best results apply in the morning before animals are turned out for pasture. Apply equivalent of 3/4 tsp. to the face or forehead of infested animals. Use a 1" paint brush to make a 6" stroke between the eyes. Apply every morning for the first two weeks and thereafter as needed. If sugar or corn syrup is not available for mixing the bait use 1 full cup of warm water.

OUTDOOR USE

Around restaurants, Drive-Ins, Ice Cream Stands, Dog Kennels, Outdoor Latrines, Compost areas -- sprinkle fly bait lightly on floors, walks, alleyways, and other horizontal surfaces where flies frequent. Repeat as necessary.

KENNELS

Treat walkways and other areas surrounding animal runs. Apply to manure as indicated for Poultry.

CAUTION

Induce vomiting if swallowed. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. KEEP OUT OF REACH OF CHILDREN, POULTRY AND DOMESTIC ANIMALS. Do not use inside homes or where milk is stored or processed. This product contains an organic phosphate which is cholinesterase inhibitor and if poisoning should occur, CALL A PHYSICIAN IMMEDIATELY. ATROPINE IS ANTIDOTAL. Harmful if swallowed, inhaled or absorbed through the skin. Destroy container when empty.

ACCEPTED WITH COMMENTS