

[193]

Front Panel

NET CONTENTS

E.P.A. Reg. No. 60023

HUB STATES 50% MALATHION
EMULSIFIABLE SPRAY

	By Weight
ACTIVE INGREDIENTS	95%
Malathion, O,O dimethyl dithiophosphate of diethyl mercaptosuccinate	50%
Xylene	45%
INERT INGREDIENTS	5%
TOTAL	100%

CAUTION ! KEEP OUT OF REACH OF CHILDREN !
SEE REVERSE PANEL FOR ADDITIONAL CAUTIONS.

ACCEPTED
JAN 3 1969
U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF PESTICIDE REGULATION
WASHINGTON, D.C. 20250
SUBJECT TO ANY OTHER COMMENTS

[293]

SIDE PANEL

DILUTION DIRECTIONS
GENERAL PEST CONTROL WORK

For general Pest control work use 2% Malathion made by diluting one part 50% Malathion Emulsifiable spray with 25 parts water.

DIRECTIONS FOR USE.

For the control of roaches, ants, spiders, silverfish, crickets, clover mites, and earwigs in and about homes, dairies, and industrial plants apply diluted solution as a coarse spray to insects with a 2% solution of Malathion. A 2% Malathion solution is made by diluting one part 50% Malathion Emulsifiable spray with 25 parts of water.

Thoroughly treat door and window casings, baseboards, and all cracks and crevices and other areas frequented by insects. Treatment of ground adjacent to buildings, foundations and lower side walls will aid in control of ants, spiders, mites, and crickets. Spot treatment can be made with the 3% Malathion however, this concentration is not recommended for general use.

THE STATES CHEMICAL AND EQUIPMENT COMPANY

200 North Hill Street, Chicago, Illinois 60601

1-31-69

5602-63 with
comments

SIDE PANEL

[393]

Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs.


Keep out of reach of children and domestic animals. Malathion upon repeated or prolonged careless use may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal.

Do not contaminate milk, milking equipment, feed or drinking water. Do not apply on animals. Remove animals and poultry from buildings before treating. Do not use in edible products area of food processing plants.

FOR FLY CONTROL IN AND AROUND

FARM BUILDINGS AND DAIRY BARN

Apply the spray solution at the rate of one gallon per 1000 square feet on unpainted surfaces where flies alight or congregate; walls, ceilings, stanchions, windows sills, fences, etc. Use a large droplet spray that wets all surfaces to the point of run off. The spray solution should not exceed a 2% concentration by weight for farm use.



With Comments

ACCEPTED WITH COMMENTS