

NITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION

VIII *Indoor*

HOMES, DAIRIES, FOOD PROCESS PLANTS

ANTS, CRICKETS, CLOVER MITES, EARWIGS, PANTRY PESTS*, SCORPIONS, SILVERFISH, SPIDERS: Use a 3% Solution of Hub States 57% Malathion made by diluting 3 tablespoonfuls of Hub States 57% Malathion liquid in 1 quart water or 6½ ounces per gallon. Apply by means of a coarse spray, paint brush or pressurized spray to window sills, baseboards, drainboards, under sinks, stoves, cracks, crevices, and to other areas frequented by insects.

BEDBUGS: Use Hub States Malathion 57% at the rate of 2 to 4 tablespoonfuls per gallon of deodorized kerosene. Apply lightly to all mattress surfaces in sufficient quantity to "mist" the fabric and generously to beds and woodwork, with special care taken to wet all possible hiding places.

CARPET BEETLES, CLOTHES MOTHS: Use a Hub States 57% Malathion 3% solution. Apply to all baseboards, floors (including areas under carpets, along margins of carpets and in closets), behind radiators and other lint accumulation areas, closet shelves and walls, and infested surface areas of carpeting. Application by means of low pressure spray or water spray will also control clothes moths in these locations.



ROACHES: Use Hub States Malathion 57% in a 3% solution and apply by means of a coarse spray, paint brush, or low pressure spray to window sills, baseboards, drainboards, under sinks, stoves and other areas frequented by insects.

IT SHOULD BE NOTED THAT WHEN USING A 3% SOLUTION INDOORS, THIS SHOULD BE LIMITED TO SPOT APPLICATIONS ONLY.

MOSQUITOES AND SMALL FLYING INSECTS: *VIII outdoors* Outdoors, use a 2% to 5% area spray, fog or aerosol. ~~As a 2% area or patio spray, dilute 58% emulsifiable liquid 1 part to 28 parts water. For 5% dilute 1-11. For mosquito larvae in standing water (Use this dosage only in intermittently flooded areas, stagnant water, temporary rain pools, sloughs and log ponds.)~~ apply Hub States 57% Malathion liquid at the rate of 13 fluid ounces (approximately ½ pound of actual Malathion) per acre. Mix in sufficient water to obtain even coverage when applied by air or ground equipment.

Lot No.

USDA Reg. No. 5602-85



Hub States Corporation

ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION
2000 North Illinois St., Indianapolis, Ind. 46202 • Phone 317 925-1

57% MALATHION

ACTIVE INGREDIENTS	By Wt.
Malathion*	57%
Aromatic Petroleum Solvent	37%
INERT INGREDIENTS	6%
	TOTAL 100%

*O,O dimethyl phosphorodithioate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions



PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED



ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION
 2000 North Illinois St., Indianapolis, Ind. 46202 • Phone 317-925-1431

57% MALATHION

ACTIVE INGREDIENTS	By Wt.
Malathion*	57%
Inert Petroleum Solvent	37%
Other INGREDIENTS	6%
TOTAL	100%

* dimethyl phosphorodithioate of diethyl mercaptosuccinate

KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions

AMOUNT OF 57% MALATHION E. C. PER 100 GALLONS WATER

ORNAMENTALS: Oyster Shell Scale*, Lace Bug—1 pt.; Euonymus Scale—1 pt. - 1½ pt.; Aphids, Mealybugs, Spider Mites, Whitefly, Four-Lined Leaf Bug, Japanese Beetle Adult, Potato Leafhopper, Tarnished Plant Bug, Thrips, Rose Leafhopper, European Pine Shoot Moth, Scurfy Scale—1½ pts.; Birch Leaf Miner, Boxwood Leaf Miner, Bagworms, Tent Caterpillar, Azalea Scale, Oak Kermes*, Pine Leaf Scale, Magnolia Scale, Fletcher Scale*, Florida Red Scale*, Juniper Scale*—2 pts.; Black Scale Crawlers—2½ pts.; Monterey Pine Scale, Soft Scale—2½ pts.; Pine Needle Scale—4 pts.; Wax Scale—2 qts., Apply in spring when crawlers are active. Repeat 1 or 2 full coverage applications at 10 day intervals.

GREENHOUSE: (In and around greenhouses and gardens.) Millipedes, Springtails, Sowbugs—Mix 1 Teaspoonful of 57% Malathion in 1 gallon of water and apply to 150 sq. ft. of soil surface or where insects congregate.

ROSES, CHRYSANTHEMUMS, CARNATIONS: Aphids, Whiteflies, Mealybugs, Thrips, Two-spotted Mite—Apply one pound of 15% aerosol per 50,000 cubic feet.

* Apply when Scale Crawlers have settled on foliage.

CAUTIONS

HARMFUL IF SWALLOWED. KEEP OUT OF REACH OF CHILDREN. Avoid contact of skin or clothing. Clothing should be washed before re-wearing. Wash thoroughly after using. Avoid prolonged or repeated contact with skin. Wear respirator when applying Malation in greenhouses or buildings. Should poisoning occur, call a physician immediately. Atropine and 2-Pam are antidotal.

Avoid contamination of food utensils, milk, milk equipment, and water. Do not use in milk processing plants. Do not use in commercial food preparation areas or in edible products areas of food processing plants.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes. Remove fish from room before applying aerosols.

Avoid applying directly to ornamental plants in homes. Keep away from heat, sparks and open flames. Destroy containers after use. Keep tightly closed while in use.

Net Contents



ACCEPTED
July 20, 1970
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND WEED KILLER ACT
 FOR ECTOPHAGOUS INSECT REGISTERED
 UNDER NO. 2112-28 SUBJECT
 TO ATTACHED COMMENTS.