

NITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANI

MIXING

Add the recommended amount of about one half the required volume of water and agitate well. When completely mixed, add the rest of the water and mix until spray emulsion is uniform.

FORAGE CROPS (ALFALFA, CLOVER, GRASS, SOYBEANS, COWPEAS, PEANUT FORAGE, AND VELVET BEANS): Apply at the following rates, using low gallonage or conventional spray equipment. Start applications at the first sign of insect infestation and repeat at 7- to 14-day intervals, as needed. Use enough water to give adequate coverage.

For Spittlebugs and Leafhoppers use 1 to 2 quarts per acre; for Alfalfa Caterpillar and Flea Beetles use 1½ to 2 quarts per acre; for Alfalfa Weevil larvae, Alfalfa Webworm, Fall Armyworm, Clover Leaf Weevil, Pea Weevil, Mexican Bean Beetle, Soybean Caterpillar, and Velvet Bean Caterpillar use 2 to 3 quarts per acre; for Cowpea Curculio and Japanese Beetle use 3 quarts per acre; for Armyworm use 3 to 4 quarts per acre.

NOTE: Apply no later than 7 days before harvest or grazing. Do not exceed 6 quarts per acre per application.

POTATOES: For Colorado Potato Beetle, Leafhoppers, and Flea Beetles use 2 quarts per acre, as needed. Do not exceed 4½ quarts per acre per application.

CATTLE (OTHER THAN DAIRY ANIMALS): As a spray for Red Short-Nosed and Long-Nosed Lice and hornflies, use 1 quart in 12½ gallons of water. Use 2 quarts of spray mixture for each average size animal. The spray should drench the entire animal until the skin is wet. If Tail Louse is present, use 2 quarts in 12½ gallons of water. Spray the entire animal, especially the switch of the tail.

On "Back Rubbers" for Horn Flies, dilute 1 gallon with 4 gallons of Diesel fuel or furnace type oil. Do Not use any other type of oil. Use one gallon of solution to saturate burlap on "Back Rubber;" soak burlap every 2 to 3 weeks.


HOGS: As a spray for Lice, use 1 quart in 12½ gallons of water. Wet the animals thoroughly with the spray mixture.

NOTE: Do Not use this material as a dip treatment. Use only as a spray.

FLIES: To help reduce populations of house and stable flies in and around farm buildings, dumps and other outdoor locations, use 1 quart in 5 gallons of water. Apply in space with mist blower, or directly to flies with hydraulic spray equipment. Do Not permit spray or spray mist to contact foodstuffs, livestock feed, watering equipment or milk handling equipment. Dairy animals should not be present while spraying. Do Not use in poultry houses.

Lot No.

USDA Reg. No. 5602-86



Hub States Corporation

ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION

METHOXYCHLOR 25

EMULSIFIABLE CONCENTRATE

FOR CERTAIN INSECT PESTS OF LIVESTOCK, FORAGE CROP POTATOES, STORED GRAIN, AND AGAINST FLIES AND ELK BARK BEETLES. CONTAINS 2 LBS. TECHNICAL METHOXYCHLOR PER GALLON.

ACTIVE INGREDIENTS	By Wt.
*Methoxychlor, Technical	25.2
Xylene	71.8
Inert Ingredients	3.0

TOTAL 100.0

*Equivalent to 22.2% 2,2-Bis (p methoxyphenyl)-1,1, 1-trichloroethane and 3% other isomers and related compounds.

DO NOT RE-USE EMPTY CONTAINER, DESTROY WHEN EMPTY

CAUTION KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions



ACCEPTED

ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED



Hub States Corporation

ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION
2000 North Illinois St., Indianapolis, Ind. 46202 - Phone 317 975 1431

METHOXYCHLOR 25-E

EMULSIFIABLE CONCENTRATE

FOR CERTAIN INSECT PESTS OF LIVESTOCK, FORAGE CROPS, POTATOES, STORED GRAIN, AND AGAINST FLIES AND ELM BARK BEETLES. CONTAINS 2 LBS. TECHNICAL METHOXYCHLOR PER GALLON.

ACTIVE INGREDIENTS	By Wt.
Methoxychlor, Technical	25.2%
Xylene	71.8%
Inert Ingredients	3.0%
	TOTAL 100.0%

Equivalent to 22.2% 2,2-Bis (p-methoxyphenyl)-1,1,1-trichloroethane and 3% other isomers and related compounds.

DO NOT RE-USE EMPTY CONTAINER, DESTROY WHEN EMPTY

CAUTION KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions

ELM BARK BEETLES: Hydraulic Sprayers: Mix 8 gallons per 100 gallons of spray. Apply enough spray to thoroughly cover all bark on trunk, limbs, and twigs (about 20 to 30 gallons of spray should cover a 50-foot elm tree).

Mist Blowers: Mix 5 gallons per 10 gallons of spray. Thoroughly cover all bark surfaces. (2 to 3 gallons of spray should cover a 50-foot elm tree.)

When to Use: Apply as a dormant treatment in the spring before leaves appear, when temperature is above 40° F. and there is no danger of freezing before spray dries.

EMPTY GRAIN BINS: To help control Rice Weevil, Granary Weevil, Lesser Grain Borer, Cadelle, Saw-toothed Grain Beetle, Confused Flour Beetle, and several other insect pests of stored grain, and help prevent infestation of newly stored grain use one gallon in 10 gallons of water. Apply during accessible or exposed stages, before grain is stored, at the rate of 2 gallons of spray per 1000 square feet. Do Not add grain to bin for at least 24 hours or until sprayed surfaces have dried thoroughly.

COMPATIBILITY

Do Not use with materials containing sulfur or other substances incompatible with oil. Do Not use on crops within 14 days of application of any material containing sulfur.

NOTE: Do Not store at temperatures below 20° F. If exposed to prolonged cold, allow to warm to 50° F. and agitate well before using. Observe the usual precautions concerning application of oil-containing products to crops during high temperatures, since many crops are then more susceptible to oil injury.

CAUTION

Do Not get on skin. In case of contact, wash with soap and water. Do Not breathe spray mist. Do Not contaminate feed or foodstuffs. Do Not use on humans. Do Not store or use near fire or open flame.

This product is toxic to fish. Keep out of any body of water. Do Not contaminate water by direct application, cleaning of equipment, or disposal of waste and containers. Apply this product only as specified on this label. Do Not contaminate water by cleaning of equipment or disposal of wastes.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Net Contents

