

◙▰◙▰◙▰◙≠®≠®≠®≠₽®₽≠®≠®≠₽®₽₽₽®

ACTIVE INGREDIENTS By Wt. O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) 2,2-Dichlorovinyl dimethyl phosphate 11,45% Total 100.00%

DANGER

1.5

**KEEP OUT OF REACH OF CHILDREN** 

(See side panels for additional cautions and antidotes.)

where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

NOTE TO PHYSICIAN: Di-Tox E insecticide is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote, 2-PAM is also antidotal and may be administered in accordance with Atropine.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes risk of any such use.

**NET CONTENTS** 

## NITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION

## DIRECTIONS FOR SPOT TREATMENT ONLY

Di-Tox E insecticide mixes readily with water or oil. Dilute the required amount of Di-Tox E in water or in a base oil such as deodorized kerosene or No. 2 fuel oil. Mix according to the following table to obtain the desired concentration of 0.25% Dursban and 0.25% DDVP Vapona or 0.5% Dursban and 0.5% DDVP Vapona insecticides. Use the higher concentrations of DDVP Vapona and Dursban to quickly reduce heavy cockroach infestations and the lower concentration of DDVP Vapona and Dursban on light infestations.

(A)Conc. Desired	WATER EMULSION Amount Required For			OIL SOLUTION Amount Required For		Mixing Ratio
			Mixing Ratio			
	1 Gal.	6 Gals.		1 Gal.	6 Gats.	
0.25% Dursban 0.25% DDVP Vapona	2-2/3 Oz.	1 Pt.	1-47	2-1/5 Oz.	13-1/5 Oz.	1-58
0.5% Dursban 0.05% DDVP Vapona	5-1/3 Oz.	1 Qt.	1-23	4-1/3 Oz.	1 Pt. 10 Oz.	1-28-5

INDOOR APPLICATION: Residential and Commercial, including restaurants, food processing plants, kennels, and horse stables. Use as a spot treatment by applying as a coarse spray or with a paint brush, to areas where cockroaches normally are found. These areas include dark corners of rooms and closets, at the base of walls, in cracks in the wall, behind baseboards, behind and beneath sinks, refrigerators, stoves and cabinets and around drainpipes. Apply in a manner to avoid spray or mist contacting food, feedstuff and water supplies. Dishes and food handling utensils should be thoroughly washed in soap and water if they have become contaminated by the application of this product. Repeat treatment as needed. NOTE: A period of 4 to 7 days after treatment is normally required for maximum effect on cockroaches.

LOT No.

าะการคาร์ที่ปรึ่งไม้ได้มีผู้ชัญญา(การ์) ก็ผู้สูญผู้ผู้มีผู้ให้เห็นก็สุดการ์ครั้งครั้งมาและเหลือมที่มีการการการก

ALC: NOT THE OWNER

USDA Reg. No. 5602-97

5

З

7

**\*** © **~** © **\*** © **\*** © **\*** © **\*** ©

