JUL 24 1974

UNDER TO ATTACHED COMMENTS.









FOR USE BY PEST CONTROL OPERATORS ONLY TO RESISTANT AND NON-RESISTANT COCKROACHES	CONTROL
ACTIVE INGREDIENTS	Bv Wt.
O,O-diethyl O-(3,5,6-trichloro-2-pyridyl)	,
phosphorothioate	12.31%
2,2-Dichlorovinyl dimethyl phosphate 11,45%	
and related compounds .86%	12,31%
Xylene.,	
INERT INGREDIENTS	5.00%
	ot 100,00%

Pest Control in Meat Packing and Food Processing Plants, Food Storage Areas and Restaurants:

NON-FOOD AREAS: Apply spot treatments to floors, walls, ceilings, or other infested areas.

FOOD AREAS: LIMITED TO (CRACK AND CREVICE TREATMENT ONLY-)
Apply a small amount of material directly into cracks and
crevices such as expansion joints between different elements
of construction or between equipment bases and the floor, wall
void, motor housing, junction boxes or switch boxes, conduits
or hollow equipment legs where the insects noted under the
directions for use hide, using a sprayer that delivers a pin
thin stream.

Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED.

Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestations. Good sanitation and proper disposal of wastes is essential in any effective insect control program.



# POTIONS I ... SPOT TREATMORT ONLY

Di-Tox E insecticide mixes readily with water or oil, Dilute the required amount of Di-Tox E in water or in a base oil such as deodorized kerosene or No. 2 fuel oil, Mix according to the following table to obtain the desired concentration of 0.25% Dursban and 0.25% DDVP Vapona or 0.5% Dursban and 0.5% DDVP Vapona insecticides. Use the higher concentrations of DDVP Vapona and Dursban to quickly reduce heavy cockroach infestations and the lower concentration of DDVP Vapona and Dursban on light infestations.

WATER EMULSION			OIL SOLUTION			
	Amount Required For		Mixing Ratio	Amount Required For		Mixing Ratio
(A)Conc. Desired	1 Gal.	6 Gals,		1 Gat.	6 Gals.	
0.25% Dursban 0.25% DDVP Vapona	2-2/3 Oz.	1 Pt.	1-47	2-1/5 O.:.	13-1/5 Oz.	1-58
0.5% Dursban 0.5% DDVP Vapona	5-1/3 Oz.	1 Qt	1-23	4-1/3 Oz.	1 Pt. 10 Oz,	1-28-5

INDOOR APPLICATION: Residential and Commercial, including restaurants, food processing plants, kennels, and horse-stables. Use as a spot treatment by applying as a coarse spray or with a paint brush, to areas where cockroaches normally are found. These areas include dark corners of rooms and closets, at the base of walls, in cracks in the wall, behind base-boards, behind and beneath sinks, refrigerators, stoves and cabinets and around drainpipes. Apply-in a menner, to avoid spray or mist contacting-food, feedstuff- and water supplies. Dishes, and food handling utensils should be thoroughly washed in coap and water if they have become contamineted by the application of this product. Repeat treatment as needed. NOTE: A period of 4 to 7 days after treatment is normally required for maximum effect on cockroaches.

LOT No.

USDA Reg. No. 5602-97

### PORTOR

Combustible liquid, do not use, spill, pour or store near heat or open flames.

Do not use in edible product areas of food processing plants.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (Keep out of smoke).

Keep out of reach of children. May be fatal if swallowed, inhaled or absorbed through skin. Do not wear contaminated clothing. May be irritating to eyes and skin. Wash thoroughly after handling. Do not breathe vapor or spray mist. Handle concentrate in ventilated area. Keep away from food, feedstuffs and water supplies. Do not let children or pets contact treated area until thoroughly dry. Flush contaminated eyes and skir, with plenty of water and for eyes get medical attention.

Do-not use-in\_edible-product areas of -food-processing plants, restaurants\_or\_other\_areas where food, is commercially-prepared or processed=Do-not use-in-serving areas while food is 'exposed.

NOTE TO PHYSICIAN: Di-Tox E insecticide is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote, 2-PAM is also antidotal and may be administered in accordance with Atropine.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes risk of any such use.

## **NET CONTENTS**

