

NITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION ADVANCED P

DIRECTIONS FOR USE

WARNING!

Harmful vapor and mist may be absorbed through skin. May produce irritation of skin and eyes, Harmful if swallowed.

Avoid breathing vapor or spray mist. Do not get in eyes or on skin. In case of skin contact, wash thoroughly with soap and water. For eyes, flush with plenty of water. Avoid contamination of feed and food stuffs.

FOR CONTROL OF TICKS AND LICE ON BEEF CATTLE, HOGS AND HORSES: Use at the rate of 1 pint in 50 gallons of water.

DIP CONTROL ON TICKS, LICE AND MANGE (SARCOPTIC AND PSOROPTIC) ON BEEF CATTLE, SHEEP, HOGS AND HORSES: Use 1 pint per 50 gallons water in dipping vat where dipping is required by state law. Water animals before dipping. Rest tired animals before treating.

Do not treat animals under 3 months of age. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

Do not dip animals within 60 days of slaughter.

Do not spray animals within 30 days of slaughter.

Do not treat dairy goats.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

"DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME."



Hub States Corporation
 ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION
 2400 North Illinois St., Indianapolis, Ind. 46202 - Phone 317/925-1431

**B-20 LINDANE
 INSECTICIDE SPRAY CONCENTRATE**

Contains

1.66 Pounds Lindane Per Gallon

Insecticide Concentrate containing 20% by weight of LINDANE to be diluted with water or deodorized Kerosene for use. Properly mixed and applied will control FLIES, MOSQUITOES, TICKS, LICE, CHIGGERS and certain other insects as listed.

ACTIVE INGREDIENTS:

Gamma isomer of Benzene Hexachloride from Lindane	20.2%
Xylene	70.7%
Diisobutylketone	4.1%
INERT INGREDIENTS	5.0%

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN!
 See Right Side Panel for Additional Cautions

FOR
 buildin
 gallon
 frame
 and o

Do Ne
 PREC
 and v
 media
 from l

Do Ne
 water.

Thoro

TICKS
 lindan
 camp
 rate c
 knaps
 most
 this s

DO N
 will g
 Keep



ACCEPTED
 4/19/72
 U.S. DEPARTMENT OF AGRICULTURE
 PESTICIDE ACT
 REGISTRATION
 No. 02-128
 SUBJECT
 COMMENTS



Hub States Corporation
 ADVANCED PRODUCTS FOR ENVIRONMENTAL SANITATION
 2400 North Illinois St., Indianapolis, Ind. 46202 - Phone 317/925-1431

**B-20 LINDANE
 INSECTICIDE SPRAY CONCENTRATE**

Contains

1.66 Pounds Lindane Per Gallon

Insecticide Concentrate containing 20% by weight of LINDANE to be diluted with water or deodorized Kerosene for use. Properly mixed and applied will control FLIES, MOSQUITOES, TICKS, LICE, CHIGGERS and certain other insects as listed.

ACTIVE INGREDIENTS:

Gamma isomer of Benzene Hexachloride from Lindane	20.2%
Xylene	70.7%
Diisobutylketone	4.1%
INERT INGREDIENTS	5.0%

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN!
 See Right Side Panel for Additional Cautions

DIRECTIONS FOR USE

FOR RESIDUAL CONTROL OF FLIES AND MOSQUITOES in barns and other farm building and in industrial plants. Mix 4 tablespoons in 1 gallon of water or 1 quart in 15 gallons of water. Apply as a coarse wet spray to walls, ceiling, beams, screens, window frames, posts, partitions and similar places where flies gather. Also to silo, manure sheds, and other adjacent structures. Repeat treatments every 30 days or as control diminishes.

CAUTION!

Do Not apply Lindane on Dairy Cattle, Dairy Barns, or in Milk Rooms.

PRECAUTIONS: Before spraying with Lindane, remove or cover all feed and cover food and water troughs or flush them out thoroughly with water from a pressure hose immediately after spraying and before spray deposit has dried. Animals should be removed from barn while Spraying.

Do Not use in poultry house. Do not contaminate milk, milking equipment, feed or drinking water.

Thoroughly air buildings after treatment before allowing persons or animals to enter.

TICKS AND CHIGGERS ON VEGETATION: A minimum dosage of 0.5 pounds of actual lindane per acre for control of ticks and chiggers on vegetation. Lawns, recreation areas, camp grounds, golf courses. Mix 1 quart thoroughly in 25 gallons of water and apply at the rate of 20 to 25 gallons per acre, or about 1 gallon to 1000 square feet. Cylindrical or knapsack-type hand sprayers may be used over small areas, but power equipment will be most satisfactory for large acreages. Treat ground litter, and the lower parts of plants with this spray.

CAUTION!

DO NOT apply Lindane where it will contaminate fruit or vegetables or where livestock will graze.
 Keep children and pets out the treated area until the spray dries.

E.P.A. Reg. #5602-120

Net Contents

