

Farmingdale Garden Labs., Inc.
136 Verdi Street
Farmingdale, New York 11735

October 14, 1971

FARMINGDALE OIL SPRAY Dormant and Summer Use

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Shake well. Use a forceful spray, wetting pests and all parts of plants thoroughly.

FOR DORMANT USE — To control Scale Insects, Red Spiders, Mites, and Mealybugs on Dormant Shrubs, Evergreens (note caution), Woody Plants, Fruit Trees (apples, pears, peaches, plums and grapes), and Shade Trees. Use 1 part Oil Spray in 25 parts of water ($\frac{1}{3}$ pt. per gal.).

GROWING SEASON — To control Scale Insects, Red Spiders, Mites, Mealybugs and White Fly Larvae on Citrus, Fruit Trees (apples, pears, peaches, plums and grapes), Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs. Use 1 part Oil Spray in 50 parts water ($\frac{1}{4}$ tablespoons per gal.). This application is also effective in the removal of sooty mold fungus through the control of white fly larvae.

AS A STICKER-SPREADER — Add 4 tablespoons to each gallon of insecticide water mixture to increase efficiency of a spray treatment, provide better contact with insects or plant surfaces and to retard weathering off of the residual materials applied.

OVICIDE: Helps to destroy insect eggs such as RED SPIDER EGGS, OAK MOTH EGGS, AND CODLING MOTH EGGS.

MADE IN U.S.A.

FARMINGDALE GARDEN LABORATORIES, INC.

136 VERDI STREET, FARMINGDALE, N.Y. 11735

E.P.A. REG. NO. 5695-10



60 SECOND LOW VISCOSITY SUPERIOR OIL

OIL SPRAY

Dormant and Summer Use

AN INSECTICIDE AND SPRAY CARRIER

ACTIVE INGREDIENTS:		by weight
Petroleum Oils		97
INERT INGREDIENTS		3
(Petroleum Oils—Minimum Unsulfonatable Residue 94%)		100

CAUTION: Keep out of reach of children.

See other warnings on side panel.

NET CONTENTS 1 QUART LIQUID

USE in Growing Season and Dormant Use Against Red Spiders, Scale Insects, Mealybugs, White Flies and certain other garden pests.

AS A PLANT POLISH — To brighten leaves of hardy house plants: Use 1 part Oil Spray in 100 parts water (3 tablespoons per gal.).

DILUTION TABLE

Tsp. — Teaspoonful	Tbs. — Tablespoonful
Add Water: 1 to 25	1 to 50
To Make: 4 Tbs.	1 qt. (32 fl. oz.)
One Gal.	4 Tsp.
Three Gal.	12 Tsp.

IF GARDEN HOSE SPRAYER IS USED — Follow directions on the bottle.

HARMFUL IF SWALLOWED

CAUTION: Wash hands after use. Do not use Oil Spray within 30 days before or after using Sulfur. Avoid spraying when temperature is excessively high or when plants are suffering from drought. Oil sprays should not be applied to broad leaved evergreens or palms when freezing or near freezing temperatures may be expected within 3 weeks after spray application. Do not use on Maidenhair Ferns, Cocos Palms, Maples, Juniper, Redbud, Sugar or Japanese Maples or Non-Woody Plants unless specified. Oil Spray may cause discoloration to Blue Spruce.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

140