

* Do Not Use on t. Augustine,
carpet, centipede or Bahia grass.

IMPORTANT

1. Be sure soil is moist and lawn healthy before treating. During drought water grass well several days in advance, and restore turf to healthy green color with fertilizer before treating.

2. Keep lawn mowed about 1½ - 2 inches during treatment.

3. Flush sprayer thoroughly with plain water after using.


4. Lawn may discolor from treatment; but, if directions are followed, turf grasses will quickly return to normal.

NOTE: FARMINGDALE CRABGRASS KILLER is a selective crabgrass killer at recommended rates of application in the presence of Kentucky, Merion Blue and Zoysia Grasses. Under ideal weather and turf conditions, such as cool days and moist turf, it is also selective in the presence of sensitive grasses, such as Fescues and Bents. Discoloration may occur in Fescues and Bentgrass areas, hence only the "Young Crabgrass" dosage is recommended when these grasses predominate. This will require a number of treatments. If spraying is delayed until crabgrass is mature in these areas.

CAUTION: Avoid contact with skin or breathing of spray mist. Avoid contamination of food or foodstuffs. Keep away from Children and domestic animals. Keep livestock off treated areas. Wash hands after using.

ANTIDOTE: If taken internally induce vomiting and call a Doctor at once.

KEEP TIGHTLY CLOSED WHEN NOT IN USE



DISODIUM METHANEARSONATE Crab Grass Killer

ACTIVE INGREDIENTS:

Disodium Methaneearsonate Anhydrous .. 12.6%

INERT INGREDIENTS .. 87.4%
100.0%

Total Arsenic as elemental,
all in water soluble form .. 5.13%

Equivalent to 20.0% Disodium
Methaneearsonate Hexahydrate

~~REGISTERED TRADE MARK~~

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE OTHER CAUTIONS ON LABEL

NET CONTENTS 1 LIQUID PINT

TREATS 2500 SQ. FT. OF LAWN

120

DIRECTIONS FOR USE

CRABGRASS CAN BE CONTROLLED BY ANY OF THE THREE METHODS BELOW. USE THE FIRST, IF YOU CAN, FOR BEST RESULTS.

1. YOUNG CRABGRASS: In early summer when crabgrass is young, apply 1 pint per 2,500 sq. ft. Follow with two more treatments at 7 day intervals. Repeated applications may be necessary if new crabgrass growth appears later in season. For smaller areas apply 4 oz. (¼ pint) per 600 sq. ft.

2. BRANCHING STAGE OF CRABGRASS: Apply 1 pint per 2,000 sq. ft. Follow with two more treatments at 7 day intervals. For smaller areas apply 1 oz. for every 125 sq. ft.

HOW TO APPLY

A. With FARMINGDALE LAWN SPRAYER which attaches to Garden Hose: First, measure 1,000 sq. ft. of lawn area. Place recommended amount of FARMINGDALE CRABGRASS KILLER in jar, then fill jar with water and spray evenly over 1,000 sq. ft. area.

B. With PRESSURE SPRAYER: Mix recommended amount of FARMINGDALE CRABGRASS KILLER as required above with 3 gallons of water and apply uniformly over the 1,000 sq. ft. area.

C. WATERING CAN. This type application is not recommended, but if it is the only means of applying, dilute required amount of FARMINGDALE CRABGRASS KILLER in 2 gallons of water and apply to 100 sq. ft., repeat this application 5 to 7 days later.

