

water again for at least 2 days. Repeat treatment in 2 or 3 weeks if necessary.

Tolerances established under the Federal Food Drug and Cosmetic Act permit the sale of crops bearing probable Sevin residues when Sevin is used in accordance with label directions.

COMPATIBILITY: This product is compatible with most commonly used fungicides and insecticides but should not be mixed with lime or other highly alkaline materials, such as Bordeaux, Lime Sulfur or Casein-Lime Stickers.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep out of reach of children and animals. Do not use on forage to be fed to dairy animals or animals being finished for slaughter. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not allow children and pets in treated areas until surfaces are dry.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from fumes.

*SEVIN IS A TRADEMARK OF UNION CARBIDE CORPORATION FOR THE ACTIVE INGREDIENT 1-NAPHTHYL N-METHYCARBAMATE

MIX THOROUGHLY AGITATE WHILE USING

MANUFACTURED FOR

Farmingdale Garden Laboratories, Inc.

FARMINGDALE, NEW YORK 11735

EPA EST. No. 904-NJ-1

E.P.A. REG. No. 5605-49

NET WT. 6 OZ.

161

ACCEPTED

APR 04 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

50% SEVIN[®] INSECT SPRAY

ACTIVE INGREDIENT

(CARBARYL 1-NAPHTHYL N-METHYCARBAMATE) _____

50%

INERT INGREDIENTS _____

50%

100%

USE ON: ROSES, ORNAMENTALS, FRUITS & VEGETABLES

ALSO KILLS: CHINCH BUGS ON LAWNS

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE OTHER WARNINGS ON SIDE PANEL

ACCEPTED

APR 04 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No.

®

50% SEVIN* INSECT SPRAY

ACTIVE INGREDIENT

(CARBARYL 1-NAPHTHYL N-METHYLCARBAMATE)

50%

INERT INGREDIENTS

50%

100%

USE ON: ROSES, ORNAMENTALS, FRUITS & VEGETABLES

ALSO KILLS: CHINCH BUGS ON LAWNS

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE OTHER WARNINGS ON SIDE PANEL

GENERAL DIRECTIONS

Mix 2 tablespoonsful of SEVIN Wettable Powder with a small amount of water to make a thin, smooth paste. Then add the paste to the remaining gallon of water (4 to 8 oz. of 50% Sevin to each 25 gallons of spray mixture). Agitate mixture well before and while spraying to assure uniformity. When insects appear direct spray to hit as many as possible and to thoroughly wet all plant surfaces. One gallon of spray mix will cover approximately 435 sq. ft. of garden area. Repeat applications at 7 to 10 day intervals or as directed for specific insects.

Sevin is not effective against mites. Where mites are a problem use a suitable miticide such as Farmingdale 50% Malathion.

FOLIAR SPRAY: For ROSES, AZALEAS, YEWs, RHODODENDRONS and other flowers and ornamental fruit and shade trees follow general directions. Make two applications at 10 day intervals, starting in early to mid-May, for leafminers on Birch, Boxwood and Holly. Hand pick large bugs at end of season and apply spray in early season for bagworm control. Spray tent caterpillar nests in the early evening.

VEGETABLE SPRAY: For control of Colorado Potato Beetle and Flea Beetle on POTATOES and Mexican Bean Beetle and Leafhoppers on BEANS. Follow general directions using 2 tablespoonsful per gallon of spray mixture. For Fruitworm, Flea Beetle and European Corn Borer on TOMATOES, EGGPLANTS and PEPPERS follow general directions using 3 to 4 tablespoonsful per gallon of water. Do not feed treated vines to meat or dairy animals. It may be applied up to and including the day of harvest on beans, tomatoes, eggplant and peppers.

FRUIT SPRAY: For control of Aphids, Coddling Moth, Apple Maggot, Red-banded leafroller on Apples and Pears (Eastern United States) and Oriental Fruit Moth, Plum Curculio and Cat-facing insects on Peaches, use 2 tablespoonsful to each gallon of spray mix applied to post bloom sprays at 10 day intervals. Do not apply within 30 days of petal fall on Apples. SEVIN may be applied up to 1 day before harvest on Apples, Pears and Peaches.

LAWN SPRAY: To control turf insects, Chinch Bug and Sod Webworms, drench each 100 sq. ft. of lawn area with 3 to 4 gallons of a spray mix containing a rounded tablespoonful per gallon. For best results apply after a rain or watering, and do not