

PLEASE FOLLOW DIRECTIONS CAREFULLY

WHEN TO SPRAY: Most lawn weeds, such as dandelion, plantain and chickweed can be sprayed successfully any time they are growing actively, spring, summer or fall. Proper fertilization after chemical weed control thickens turf and prevents reinfestation.

Spray when the air is still, on a calm day, early morning or evening to avoid spray drift which can damage desirable plants.

SAFE ON TURF GRASSES: Most lawn grasses, including bluegrasses, fescues, Bermudas, and Zoysias will not be injured by proper application. To control weeds in bentgrass, St. Augustine grass or newly seeded lawns, use half the rate recommended below. Delay treating newly seeded lawns until they have been mowed at least twice. Lawns can be reseeded three weeks after spraying if there has been a good rain or deep watering.

Do not use in flower, vegetable or ornamental plantings.

AMOUNTS TO USE: Measure off area to be treated. Apply one ounce of Lawn Weed Killer in a gallon of water for each 300 sq. ft. of lawn. Use a watering can. For larger areas, use Farmingdale Lawn Sprayer. Place 8 ozs. of Farmingdale Lawn Weed Killer in jar, fill with water to 8 gal. mark. Spray contents on 2400 sq. ft. of lawn.

FOR STUMP KILLING: Use one pint in one quart of diesel oil or kerosene and apply spray to all areas of the stump to prevent new growth.

POISON IVY CONTROL: Apply 8 oz. in 2 1/2 gallons of water. Thoroughly wet all leaves and stems with a drenching spray.

Do not use same sprayer for other weed killing purposes. When diluting, mix thoroughly before using. If mixture is allowed to stand for any length of time before using, it should be stirred again.



CONTAINS SILVEX ☆

2,4-D PLUS 2,4,5-TP ☆
**LAWN WEED &
DANDELION KILLER**

• **KILLS POISON IVY** • **KILLS TREE STUMPS**

KILLS: Dandelion, Chickweed, Clover, Plantain and many Broadleaf Weeds

NET CONTENTS 1 PINT LIQUID **168**
TREATS 4800 SQUARE FEET

WARNING: KEEP OUT OF REACH OF CHILDREN.
READ OTHER WARNINGS ON SIDE PANEL.

5605-88
RECORDED
JAN 10 1968
FARMINGDALE GARDEN LABORATORIES, INC.

ACTIVE INGREDIENTS:

Ethylhexyl Ester of 2,4-Dichlorophenoxyacetic Acid.....15.2%
Equivalent to 10.0% 2,4-Dichlorophenoxyacetic Acid;
Isooctyl Ester of Silvex 2-(2,4,5-Trichlorophenoxy Propionic Acid).....7.1%
Equivalent to 5.0% Silvex [2-(2,4,5-Trichlorophenoxy propionic acid.)

INERT INGREDIENTS:77.7%
100.0%

WARNING: Avoid spray drift to susceptible plants as this product may injure ornamental shrubs, trees, flowers and vegetables, etc. (Coarse sprays are less likely to drift.) Do not use the same spray equipment for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Under very high temperature of use, vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation purposes. Avoid contact with skin, eyes or clothing. Do not use in a greenhouse. Keep out of reach of children.

KEEP CONTAINER CLOSED
STORE AWAY FROM HEAT
AND OPEN FLAME

Manufactured For
FARMINGDALE GARDEN LABORATORIES, INC.
136 VERDI STREET, FARMINGDALE, N. Y.

U.S.D.A. REG. NO. 5605-88