


FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York



11-22-66

FARMINGDALE LAWN & GARDEN FUNGICIDE CONTAINING MANEB with Zinc

Front Panel:

FARMINGDALE LAWN & GARDEN FUNGICIDE
Containing Maneb with Zinc

Controls Many Common Diseases on Turf, Ornamentals and Vegetables

ACTIVE INGREDIENT:

Maneb (Manganese ethylene bisdithiocarbamate) 80.0%

(metallic manganese equivalent-16.5%)

INERT INGREDIENT:

(including zinc) 20.0%
100.0%

U.S. Patents 2,504,404; 2,974,156
and Patent Pending

CAUTION: KEEP OUT OF REACH OF CHILDREN

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes get medical attention.

NET WEIGHT: ONE POUND
U.S.D.A. Reg.No. 5605-

Mfg. for: FARMINGDALE GARDEN LABS., INC.
FARMINGDALE, NEW YORK

GENERAL DIRECTIONS: Mix with a small amount of water to make a thin paste, then add to balance of water. Stir well and thoroughly spray all plant surfaces. This fungicide is believed to be compatible with most commonly used pesticides.

TURF GRASSES-For stem, leaf and stripe rusts, use 2 tablespoons per gallon. Apply 2 gallons per 1000 sq.ft. when rust pustules are first seen and repeat every 7 to 14 days as long as rust threatens. For Sclerotinia dollar spot, use 6 to 8 oz. per 1000 sq.ft. in sufficient water to provide adequate coverage. Start application at first sign of disease and repeat every 7 to 14 days until disease threat is past.

For Melting Out, use 3 to 4 oz. per 1000 sq.ft. in sufficient water to provide adequate coverage. Start application at first sign of disease and repeat every 7 to 14 days until disease threat is past.

For control of Rhizoctonia brown patch, use at 3 oz. per 1000 sq.ft. in sufficient water to provide adequate coverage. Start application at first sign of disease and repeat every 5 to 10 days until disease threat is past. Follow local recommendations for further details on timing.

ORNAMENTALS-Use 1½ level tablespoons per gallon in full coverage spray. Begin spraying when plants are well leafed-out and apply at weekly intervals throughout the season or follow local commercial florist recommendations for control of: ROSE-black spot, DAHLIA-Botrytis, PANSY-Anthraco-nose, ZINNIA-Leaf blight, GLADIOLUS-Stemphyllium, Curvularia and Botrytis, CARNATION-Leaf Spot, LILY-Botrytis, SNAPDRAGON-Rust, PEONY-Phytophthora and Botrytis blights. Apply in early Spring and early Fall, drenching soil surrounding plant as well as foliage. Destroy all infected parts promptly. FLOWERJNG DOGWOOD-Anthraco-nose. Apply when buds begin to open and again when bracts have fallen, and 4 weeks later and again in late summer after flower buds for next season are formed. VEGETABLE CROPS--Use 1½ tablespoonfuls per gallon of water. Apply at the rate of 1 gallon to 450 sq.ft.

FARMINGDALE GARDEN LABS., INC. Page 2
136 Verdi Street
Farmingdale, New York

11-22-66

FARMINGDALE LAWN & GARDEN FUNGICIDE
(continued)

CONTAINING MANEB with Zinc

TOMATOES-Early and Late Blight, Anthracnose, Stemphyllium (gray leaf Spot), Septoria leaf spot and Cladosporium gray mold on tomatoes in field, greenhouse or plant beds. Begin spraying when disease is first reported in area and repeat at 7 to 10 day intervals or oftener during severe late blight conditions. To avoid damage do not use on tender young plants under glass. CUCUMBERS AND SQUASH-Downy mildew, anthracnose and Pythium fruit rot. Begin when runners form and repeat weekly as needed. CARROTS-Leaf spot diseases. Start applications when plants are 6 weeks old or when disease threatens. Repeat at 7 to 10 day intervals. BEANS-(Green and dry)-Downy mildew, rust. Apply when disease first appears and repeat at 7 day intervals. Do not apply to green beans within 4 days of harvest. EGGPLANT and PEPPERS-Fruit rot, anthracnose. Apply as fruits begin to form and repeat at weekly intervals through harvest.

CAUTION: May cause irritations of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes get medical attention.

IMPORTANT: Keep away from fires and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of this product and may also generate flammable vapors. Keep container closed when not in use.

STATEMENT OF INERT INGREDIENTS

This material is Dithane M-22 Special

formulated by Rohm & Haas of Philadelphia, Pa.

All label claims are based on Rohm & Haas Dithane labels.