

FARMINGDALE GARDEN LABS., INC.
136 Vondra Street
Farmingdale, New York

11-27-66

FARMINGDALE DANDELION & CHICKWEED KILLER with BANVEL D
For Broadleaf Turf Weed Control on Lawns and Golf Courses.

ACTIVE INGREDIENTS

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)*	3.43%
Dimethylamine Salts of Related Acid:	0.55%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid**	17.27%
INERT INGREDIENTS	78.80%
	100.00%

*Equivalent to 2.84% 3,6-dichloro-o-anisic acid (dicamba)
or 0.25 pounds per gallon.

**Equivalent to 14.27% 2,4-Dichlorophenoxyacetic acid or 1.25 pounds per gallon

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panel for additional cautions

NET CONTENTS:
PINT

Mfg. for FARMINGDALE GARDEN LABS., INC.
FARMINGDALE, NEW YORK

U.S.D.A. REG. NO. 5605-

Camcraman's Note

Poor Copy.

DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Hambit, Wild Onion and many other broadleaf weeds in the Spring...**FOLLOW DILUTION IN SCHEDULE A.** For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Quail, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist, repeat application in 2 to 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

DILUTION CHART

<u>SCHEDULE A</u>	<u>SCHEDULE B</u>	<u>WATER GALLONS</u>	<u>AREA SQ. FT.</u>
1 Tablespoonful 1/2 fl. oz.	3.5 Teaspoonfuls 0.6 fl. oz.	1	200
5 Tablespoonfuls 2 1/2 fl. oz.	6 Tablespoonfuls 3 fl. oz.	5	1000
1 pint	19 fl. oz.	30	6500

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS: To kill Poison Ivy, Rag Weed, Morning-Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if regrowth develops. Use 1.5 fl. oz. in 1 gallon of water or 1 cupful in 5 gallons of water.

ACCEPTED

FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York

11-22-66

FARMINGDALE DANDELION & CHICKWEED KILLER WITH BANVEL D

(Continued)

CUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or St. Augustine nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

Contamination of shallow fish-bearing water at application rates recommended on this label may kill fish.

Camerman's Note

Poor Copy

11-22-66
Sent 11/25

FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York

BEST DOCUMENT AVAILABLE

11-27-66

FARMINGDALE DANDELION & CHICKWEED KILLER
For Broadleaf Turf Weed Control on Lawns and Golf Courses.

ACTIVE INGREDIENTS

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)*	3.43%
Dimethylamine Salts of Related Acids	0.50%
Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid**	17.07%
INERT INGREDIENTS	79.00%
	<u>100.00%</u>

*Equivalent to 2.04% 3,6-dichloro-o-anisic acid (dicamba)
or 0.25 pounds per gallon.

**Equivalent to 14.27% 2,4-dichlorophenoxyacetic acid or 1.25 pounds per gallon
CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panel for additional cautions

NET CONTENTS:
PINT

Mfg. for FARMINGDALE GARDEN LABS., INC.
FARMINGDALE, N. Y.

U.S.D.A. REG. NO. 5605-

DIRECTIONS:

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Ragwort, Wild Onion and many other broadleaf weeds in the Spring...**FOLLOW DILUTION IN SCHEDULE A.** For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Quail, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist, repeat application in 2 to 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

DILUTION CHART

<u>SCHEDULE A</u>	<u>SCHEDULE B</u>	<u>WATER GALLONS</u>	<u>AREA Sq. Ft.</u>
1 Tablespoonful 1/2 fl. oz.	3.5 Teaspoonfuls 0.6 fl. oz.	1	200
5 Tablespoonfuls 2 1/2 fl. oz.	6 Tablespoonfuls 3 fl. oz.	5	1000
1 pint	19 fl. oz.	30	6500

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS: To kill Poison Ivy, Rag Weed, Morning-Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if regrowth develops. Use 1.5 fl. oz. in 1 gallon of water or 1 cupful in 5 gallons of water.

ACCEPTED

5605
-95

FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York

11-22-66

FARMINGDALE DANDELION & CHICKWEED KILLER WITH BANVEL D[®]

(Continued)

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

Contamination of shallow fish-bearing water at application rates recommended on this label may kill fish.

RECEIVED
NOV 22 1966
C-17-1967
5605-95

5605-95

FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York

11-22-66

BEST DOCUMENT AVAILABLE

CHART BASED ON FOLLOWING:

SCHEDULE A

1 pint/6500 sq.ft. - 1 gal. to 52,000 sq. ft. or 0.03 gals. per acre.

1 fl. oz. to 409.5 sq.ft.

1 tablespoon to 205 sq.ft.

5 Tablespoons to 1000 sq.ft.

1.05 lbs. 2,4-D Acid equivalent and 0.21 lbs. Dicamba/acre

SCHEDULE B

~~XXXXXXXXXXXXXXXXXXXX~~

- pint/5645 sq.ft.

1 gal/acre

1 fl. oz. to 360 sq.ft.

0.5 fl. oz. to 204 sq.ft.

3.5 Tablespoons to 200 sq.ft.

2.94 fl. oz. to 1000 sq.ft.

6 Tablespoons to 1000 sq.ft.

19.1 fl. oz. to 6500 sq.ft.

1.25 lbs. 2,4-D Acid equivalent and

0.25 lbs. Dicamba/acre

If SCHEDULE B is repeated in a few weeks

Total 2.50 lbs. 2,4-D acid and 0.5 lbs. of Dicamba.

5605-95

INERT INGREDIENT STATEMENT

Water 78.80%

BEST DOCUMENT AVAILABLE

11-77-66

FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York

FARMINGDALE DANDELION & CHICKWEED KILLER WITH BAIVEL D 8
For Broadleaf Turf Weed Control on Lawns and Golf Courses.

ACTIVE INGREDIENTS

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)* 3.43%
Dimethylamine Salts of Related Acids 0.55%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid** 17.22%

INERT INGREDIENTS

78.80%
100.00%

*Equivalent to 2.84% 3,6-dichloro-o-anisic acid (dicamba)
or 0.25 pounds per gallon.

**Equivalent to 14.27% 2,4-Dichlorophenoxyacetic acid or 1.25 pounds per gallon

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panel for additional cautions

NET CONTENTS:
PINT

REG. for FARMINGDALE GARDEN LABS. INC.
FARMINGDALE, NEW YORK

U.S.D.A. REG. NO. 3605-

DIRECTIONS

~~LAWN WEEDS:~~ For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Hepbit, Wild Onion and many other broadleaf weeds in the Spring...
DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Galia, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist, repeat application in 2 to 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

DILUTION CHART

SCHEDULE A	SCHEDULE B	WATER GALLONS	AREA SQ. FT.
1 Tablespoonful 1 fl. oz.	3.5 Teaspoonfuls 0.6 fl. oz.	1	200
5 Tablespoonfuls 2 1/2 fl. oz.	6 Tablespoonfuls 3 fl. oz.	5	1000
1 pint	19 fl. oz.	30	6500

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS: To kill Poison Ivy, Rag Weed, Morning-Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if regrowth develops. Use 1.5 fl. oz. in 1 gallon of water or 1 cupful in 5 gallons of water.

ACCEPTED

5605
-98

FARMINGDALE GARDEN LABS., INC.
136 Verdi Street
Farmingdale, New York

BEST DOCUMENT AVAILABLE

11-22-66

FARMINGDALE DANDELION & CHICKWEED KILLER WITH BANVEL D @

(Continued)

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of the reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra, nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

Contamination of shallow fish-bearing water at application rates recommended on this label may kill fish.

APPROVED
APR 17 1967
5605-95

5605-95

FAIRHURD GARDEN LABS., INC.
136 Verdi Street
Farmlingdale, New York

11-22-66

BEST DOCUMENT AVAILABLE

CHART BASED ON FOLLOWING:

SCHEDULE A

1 pint/6500 sq.ft. - 1 gal. to 52,000 sq. ft. or 0.833 gals. per acre.

1 fl. oz. to 409.5 sq.ft.
Tablespoon to 205 sq.ft.
5 Tablespoonfuls to 1000 sq.ft.

1.05 lbs. 2,4-D Acid equivalent and 0.21 lbs. Dicamba/acre

SCHEDULE B

~~EXPERIMENTAL~~

1 pint/5445 sq.ft. 1 gal/acre

1 fl. oz. to 940 sq.ft.
0.5 fl. oz. to 204 sq.ft.
3.5 Teaspoonfuls to 200 sq.ft.
2.94 fl. oz. to 1000 sq.ft.
6 Tablespoonfuls to 1000 sq.ft.
19.1 fl. oz. to 6500 sq.ft.

5605-95

1.25 lbs. 2,4-D Acid equivalent and
0.25 lbs. Dicamba/acre

If SCHEDULE B is repeated in a few weeks
Total 2.50 lbs. 2,4-D acid and 0.5 lbs. of Dicamba.

NETT INGREDIENT STATEMENT

Water 78.80%