

FARMINGDALE GARDEN LABS., INC.  
136 Vondt Street  
Farmingdale, New York

11-22-66

CHART BASED ON FOLLOWING:

**SCHEDULE A**

1 pint/6500 sq.ft. - 1 gal. to 52,000 sq. ft. or 0.838 gals. per acre.

1 fl. oz. to 409.5 sq.ft.

1 Tablespoon to 205 sq.ft.

5 Tablespoons to 1000 sq.ft.

1.05 lbs. 2,4-D Acid equivalent and 0.21 lbs. Dicamba/acre

**SCHEDULE B**

**CONCENTRATION TABLE**

1 pint . . . . . 1 gal/acre

1 . . . . . ft.  
1 . . . . . sq.ft.  
1 . . . . . to 200 sq.ft.  
1 . . . . . 1000 sq.ft.  
1 . . . . . to 1000 sq.ft.  
1 . . . . . 5500 sq.ft.

1.25 lbs. 2,4-D Acid equivalent and  
0.25 lbs. Dicamba/acre

If SCHEDULE B is repeated in a few weeks  
Total 2.50 lbs. 2,4-D acid and 0.5 lbs. of Dicamba.

Chemical Note

Poor Copy

**NET INGREDIENT STATEMENT**

Water . . . . . 78.30%

8-22-67


FARMINGDALE GARDEN LABS., INC.  
136 Verdi Street  
Farmingdale, L.I.

FARMINGDALE LAWN WEED AND DANDELION KILLER - Contains Banvel D F

Also Controls Chickweed and Certain Broadleaf Lawn Weeds  
POISON IVY KILLER

CAUTION: KEEP OUT OF REACH OF CHILDREN  
See side panel for additional cautions.

Net Contents: 1 Liquid Quart Treats 8600 Square Feet  
Made For: FARMINGDALE GARDEN LABS., INC.  
FARMINGDALE, N.Y.



DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, ~~eg~~ Chickweed, Clover, Plantain, Docks, Knotweed, Henbit, Wild Onion and many other broadleaf weeds in the Spring... FOLLOW DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Oxalis, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist, repeat application in 2 or 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

<u>DILUTION CHART</u>		<u>Water Gallons</u>	<u>Area</u> <u>Sq. Ft.</u>
<u>Schedule A</u>	<u>Schedule B</u>		
4 teaspoonfuls	5 teaspoonfuls	1	200
7 tablespoonfuls	8 tablespoonfuls	5	1000
1 Pint	19 Oz.	20	4300

ACTIVE INGREDIENTS:

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)*	2.28%
Dimethylamine Salts of Related Acids	0.37%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid**	12.0%
<u>INERT INGREDIENTS:</u>	<u>85.35%</u>
	100.00%

\*Equivalent to 1.88% 3,6-dichloro-o-anisic acid (dicamba) or 0.17 pounds per gallon.

\*\*Equivalent to 10% 2,4-Dichlorophenoxyacetic acid or 0.88 pounds per gallon.

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS:

To kill Poison Ivy, Rag Weed, Morning Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if re-growth develops. Use 2 fl.oz. in each gallon of water. ~~xx~~

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable tree or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds,

8-22-67

FARMINGDALE GARDEN LABS., INC.  
136 Verdi Street  
Farmingdale, L.I.

FARMINGDALE LAWN WEED AND DANDELION KILLER - Contains Banvel D F

Also Controls Chickweed and Certain Broadleaf Lawn Weeds  
POISON IVY KILLER

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for additional cautions.

Net Contents: 1 Liquid Quart Treats 8600 Square Feet  
Made For: FARMINGDALE GARDEN LABS., INC.  
FARMINGDALE, N.Y.

ACCEPTED

OCT 3

RECEIVED  
FARMINGDALE GARDEN LABS., INC.  
FARMINGDALE, N.Y.

5605-99

DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Hepbit, Wild Onion and many other broadleaf weeds in the Spring... FOLLOW DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Oxalis, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist, repeat application in 2 or 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

<u>DILUTION CHART</u>		<u>Water Gallons</u>	<u>Area Sq.Ft.</u>
<u>Schedule A</u>	<u>Schedule B</u>		
4 teaspoonfuls	5 teaspoonfuls	1	200
7 tablespoonfuls	8 tablespoonfuls	5	1000
1 Pint	19 Oz.	20	4300

A. VE INGREDIENTS:

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)\* 2.28%

Dimethylamine Salts of Related Acids 0.37%

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid\*\* 12.0%

INERT INGREDIENTS: 85.35%

100.00%

\*Equivalent to 1.88% 3,6-dichloro-o-anisic acid (dicamba) or 0.17 pounds per gallon.

\*\*Equivalent to 10% 2,4-Dichlorophenoxyacetic acid or 0.88 pounds per gallon.

WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS:

To kill Poison Ivy, Rag Weed, Morning Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if re-growth develops. Use 2 fl.oz. in each gallon of water.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable tree or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds,

BEST DOCUMENT AVAILABLE

FARMINGDALE GARDEN LABS., INC.

-2-


8-22-67

FARMINGDALE LAWN WEED & DANDELION KILLER - Contains Banvel D

insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is ~~not~~ empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf. Contamination of shallow fish-bearing water at application rates recommended on this label may kill fish.

U.S.D.A. Registration No. 5605-

5605-99



8-22-67


FARMINGDALE GARDEN LABS., INC.  
136 Verdi Street  
Farmingdale, L.I.

FARMINGDALE LAWN WEED AND DANDELION KILLER - Contains Banvel D r

Also Controls Chickweed and Certain Broadleaf Lawn Weeds  
POISON IVY KILLER

CAUTION: KEEP OUT OF REACH OF CHILDREN  
See side panel for additional cautions.

Net Contents: 1 Liquid Quart Treats 8600 Square Feet  
Made For: FARMINGDALE GARDEN LABS., INC.  
FARMINGDALE, N.Y.



#### DIRECTIONS

LAWN WEEDS: For the most effective control at the lowest dosage, apply when the weeds are growing vigorously after a Spring rain. Carefully measure the area to be treated and apply the correct dosage evenly over the entire area. Use either a tank type or hose-end sprayer adjusted to deliver a coarse spray. For Dandelions, Chickweed, Clover, Plantain, Docks, Knotweed, Henbit, Wild Onion and many other broadleaf weeds in the Spring... FOLLOW DILUTION IN SCHEDULE A. For late summer and fall applications and for hard-to-kill weeds such as Sheep Sorrel, Oxalis, Yarrow and mature Knotweed, increase the dosages to Schedule B. If weeds persist, repeat application in 2 or 3 weeks. Wild Onions will require repeated applications in late fall and early spring.

<u>DILUTION CHART</u>		<u>Water Gallons</u> <del>Schedule B</del>	<u>Area</u> <u>Sq. Ft.</u>
<u>Schedule A</u>	<u>Schedule B</u>		
4 teaspoonfuls	5 teaspoonfuls	1	200
7 tablespoonfuls	8 tablespoonfuls	5	1000
1 Pint	19 Oz.	20	4300

#### ACTIVE INGREDIENTS:

Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)*	2.28%
Dimethylamine Salts of Related Acids	0.37%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid**	12.0%
INERT INGREDIENTS:	85.35%
	100.00%

\*Equivalent to 1.88% 3,6-dichloro-o-anisic acid (dicamba) or 0.17 pounds per gallon.

\*\*Equivalent to 10% 2,4-Dichlorophenoxyacetic acid or 0.88 pounds per gallon.

#### WEEDS IN VACANT LOTS, FENCE ROWS AND SIMILAR AREAS:

To kill Poison Ivy, Rag Weed, Morning Glory, Thistles and many other broadleaf weeds, spray when the weeds are actively growing. Wet all plant surfaces. Repeat spray if re-growth develops. Use 2 fl.oz. in each gallon of water.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact flush with abundance of water; for eyes, get medical attention. Do not take internally. Keep out of reach of children.

IMPORTANT: Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees and other desirable crop plants. Do not treat under drip line or to the root systems of sensitive ornamental shrubs and plants. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable tree or shrubs or where it may contaminate water. Do not store in direct contact with fertilizer, seeds,

BEST DOCUMENT AVAILABLE

FARMINGDALE GARDEN LABS., INC.

-2-

8-22-67

14

FARMINGDALE LAWN WEED & DANDELION KILLER - Contains Banvel D

insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is ~~not~~ empty, immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf. Contamination of shallow fish-bearing water at application rates recommended on this label may kill fish.

U.S.D.A. Registration No. 5605-

5605-99

